
SANTA CRUZ BIOTECHNOLOGY, INC.

Sall2 (K-16): sc-27120



BACKGROUND

Sall1 (SALL1, sal-like 1, TBS, HSAL1) and Sall2 (SALL2, sal-like 2, HSAL2, p150 (Sal2)) are mammalian homologs of the *Drosophila* region-specific home-otic gene spalt (sal), which encodes a zinc finger-containing transcription regulator. *Drosophila* spalt (sal) is an essential genetic component required for the specification of posterior head and anterior tail as opposed to trunk. Mammalian Sall1 may mediate higher order chromatin structure, and may be a component of a distinct heterochromatin-dependent silencing process. Sall1 is present in kidney, brain and liver. Sall2 is a p53-independent regulator of p21 and BAX, and can function in some cell types as a regulator of cell growth and survival. Human Sall2 is present in brain, heart, kidney or pancreas. Sall1 and Sall2 are expressed in different areas of the fetal brain that may represent distinct sets of neurons.

REFERENCES

- Kohlhase, J., Schuh, R., Dowe, G., Kuhnlein, R.P., Jackle, H., Schroeder, B., Schulz-Schaeffer, W., Kretzschmar, H.A., Kohler, A., Muller, U., Raab-Vetter, M., Burkhardt, E., Engel, W., Stick, R. 1996. Isolation, characterization, and organ-specific expression of two novel human zinc finger genes related to the *Drosophila* gene spalt. Genomics 38: 291-298.
- Ma, Y., Li, D., Chai, L., Luciani, A.M., Ford, D., Morgan, J., Maizel, A.L. 2001. Cloning and characterization of two promoters for the human HSAL2 gene and their transcriptional repression by the Wilms' tumor suppressor gene product. J. Biol. Chem. 276: 48223-48230.
- Nielsen, T.O., Hsu, F.D., O'Connell, J.X., Gilks, C.B., Sorensen, P.H., Linn, S., West, R.B., Liu, C.L., Botstein, D., Brown, P.O., van de Rijn, M. 2003. Tissue microarray validation of epidermal growth factor receptor and Sall2 in synovial sarcoma with comparison to tumors of similar histology. Am. J. Pathol. 163: 1449-1456.
- Sato, A., Matsumoto, Y., Koide, U., Kataoka, Y., Yoshida, N., Yokota, T., Asashima, M., Nishinakamura, R. 2003. Zinc finger protein Sall2 is not essential for embryonic and kidney development. Mol. Cell Biol. 23: 62-69.
- Li, D., Tian, Y., Ma, Y., Benjamin, T. 2004. p150(Sal2) is a p53-independent regulator of p21(WAF1/CIP). Mol. Cell Biol. 24: 3885-3893.
- SWISS-PROT/TrEMBL (Q9Y467). World Wide Web URL: http://www. expasy.ch/sprot/sprot-top.html

SOURCE

Sall2 (K-16) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of Sall2 of mouse origin.

PRODUCT

Each vial contains 200 μ g lgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-27120 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

Available as TransCruz reagent for Gel Supershift and ChIP applications, sc-27120 X, 200 $\mu g/0.1$ ml.

APPLICATIONS

Sall2 (K-16) is recommended for detection of Sall2 of mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Sall2 (K-16) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker[™] compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker[™] Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluo-rescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz^{™™} Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.