

# C1QBP (B-6): sc-271201

## BACKGROUND

The human complement subcomponent C1q associates with C1r and C1s in order to yield the first component of the serum complement system (SCS). The SCS contains over 30 glycoproteins that influence physiological mechanisms of the body in response to immune complex (the classical pathway), carbohydrate (the lectin pathway) or bacterial (alternative pathway) initiation. C1q binding protein (C1QBP), also designated gC1q-R, p32 (p33) or HABP1 (hyaluronan-binding protein 1), is known to bind the globular heads of C1q molecules and inhibit C1 activation. C1QBP has been described as a complement receptor for C1q on B cells, neutrophils and mast cells. The C1QBP protein may form homodimers. C1QBP is expressed in vascular endothelial cells and has been found to be a multifunctional protein interacting with elements of complement, coagulation and kinin systems. In addition, C1QBP is a subunit of pre-mRNA splicing factor SF2/ASF.

## CHROMOSOMAL LOCATION

Genetic locus: C1QBP (human) mapping to 17p13.2; C1qbp (mouse) mapping to 11 B4.

## SOURCE

C1QBP (B-6) is a mouse monoclonal antibody raised against amino acids 1-282 representing full length C1QBP of human origin.

## PRODUCT

Each vial contains 200 µg IgG<sub>1</sub> kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## APPLICATIONS

C1QBP (B-6) is recommended for detection of C1QBP of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for C1QBP siRNA (h): sc-42880, C1QBP siRNA (m): sc-42881, C1QBP shRNA Plasmid (h): sc-42880-SH, C1QBP shRNA Plasmid (m): sc-42881-SH, C1QBP shRNA (h) Lentiviral Particles: sc-42880-V and C1QBP shRNA (m) Lentiviral Particles: sc-42881-V.


Molecular Weight of C1QBP: 33 kDa.

Positive Controls: K-562 whole cell lysate: sc-2203, MCF7 whole cell lysate: sc-2206 or NIH/3T3 whole cell lysate: sc-2210.


## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgGκ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

## DATA



C1QBP (B-6): sc-271201. Western blot analysis of C1QBP expression in MCF7 (A), K-562 (B), C2C12 (C), NIH/3T3 (D), L8 (E) and RAT2 (F) whole cell lysates.



C1QBP (B-6): sc-271201. Immunofluorescence staining of formalin-fixed Hep G2 cells showing cytoplasmic localization (A). Immunoperoxidase staining of formalin fixed, paraffin-embedded human duodenum tissue showing cytoplasmic staining of glandular cells (B).

## SELECT PRODUCT CITATIONS

- Ounap, K., et al. 2015. The stability of ribosome biogenesis factor WBSR22 is regulated by interaction with TRMT112 via ubiquitin-proteasome pathway. *PLoS ONE* 10: e0133841.
- Liu, Z., et al. 2015. The roles of p38 MAPK and ERK1/2 in coplanar polychlorinated biphenyls-induced apoptosis of human extravillous cytotrophoblast-derived transformed cells. *Cell. Physiol. Biochem.* 36: 2418-2432.
- Liang, X.H., et al. 2017. RNase H1-dependent antisense oligonucleotides are robustly active in directing RNA cleavage in both the cytoplasm and the nucleus. *Mol. Ther.* 25: 2075-2092.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.