

LETM1 (D-5): sc-271235

BACKGROUND

LETM1 (leucine zipper-EF-hand-containing transmembrane protein 1, mitochondrial) is a 739 amino acid protein that localizes to the mitochondrial membrane and contains one LETM1 domain and two EF-hand calcium-binding domains. Expressed in all fetal and adult tissues, LETM1 has a leucine zipper motif, a transmembrane domain and several phosphorylation sites and, via its EF-hand domains, may function as a calcium-binding protein. Additionally, LETM1 is thought to be involved in maintaining normal mitochondrial function and overall cell viability. Human LETM1 shares 84% similarity with its mouse counterpart, suggesting a conserved role between species. Deletions in the gene encoding LETM1 are associated with Wolf-Hirschhorn syndrome (WHS), a congenital syndrome characterized by a number of abnormalities, including mental retardation, seizures, heart defects, fused teeth, hearing loss, a webbed neck and renal abnormalities.

REFERENCES

- Endele, S., et al. 1999. LETM1, a novel gene encoding a putative EF-hand Ca^{2+} -binding protein, flanks the Wolf-Hirschhorn syndrome (WHS) critical region and is deleted in most WHS patients. *Genomics* 60: 218-225.
- Rauch, A., et al. 2001. First known microdeletion within the Wolf-Hirschhorn syndrome critical region refines genotype-phenotype correlation. *Am. J. Med. Genet.* 99: 338-342.
- Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 604407. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Zollino, M., et al. 2003. Mapping the Wolf-Hirschhorn syndrome phenotype outside the currently accepted WHS critical region and defining a new critical region, WHSCR-2. *Am. J. Hum. Genet.* 72: 590-597.

CHROMOSOMAL LOCATION

Genetic locus: LETM1 (human) mapping to 4p16.3; Letm1 (mouse) mapping to 5 B2.

SOURCE

LETM1 (D-5) is a mouse monoclonal antibody raised against amino acids 440-739 mapping at the C-terminus of LETM1 of human origin.

PRODUCT

Each vial contains 200 µg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

LETM1 (D-5) is available conjugated to agarose (sc-271235 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-271235 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-271235 PE), fluorescein (sc-271235 FITC), Alexa Fluor® 488 (sc-271235 AF488), Alexa Fluor® 546 (sc-271235 AF546), Alexa Fluor® 594 (sc-271235 AF594) or Alexa Fluor® 647 (sc-271235 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-271235 AF680) or Alexa Fluor® 790 (sc-271235 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA

APPLICATIONS

LETM1 (D-5) is recommended for detection of LETM1 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for LETM1 siRNA (h): sc-89079, LETM1 siRNA (m): sc-146712, LETM1 shRNA Plasmid (h): sc-89079-SH, LETM1 shRNA Plasmid (m): sc-146712-SH, LETM1 shRNA (h) Lentiviral Particles: sc-89079-V and LETM1 shRNA (m) Lentiviral Particles: sc-146712-V.

Molecular Weight of LETM1: 85 kDa.


Positive Controls: HeLa whole cell lysate: sc-2200, A549 whole cell lysate: sc-2413 or Jurkat whole cell lysate: sc-2204.

RECOMMENDED SUPPORT REAGENTS


To ensure optimal results, the following support reagents are recommended:

- 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.
- 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).
- 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.
- 4) Immunohistochemistry: use m-IgGκ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

DATA



LETM1 (D-5): sc-271235. Western blot analysis of LETM1 expression in HeLa (A), F9 (B), A549 (C), Jurkat (D), MOLT-4 (E) and Ramos (F) whole cell lysates.



LETM1 (D-5): sc-271235. Immunofluorescence staining of methanol-fixed HeLa cells showing cytoplasmic localization (A). Immunoperoxidase staining of formalin fixed, paraffin-embedded human small intestine tissue showing cytoplasmic staining of glandular cells (B).

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.