

TCP-1 η (C-6): sc-271439

BACKGROUND

The protein TCP-1 (t complex polypeptide 1) is a subunit of the hetero-oligomeric complex CCT (chaperonin containing TCP-1) present in the eukaryotic cytosol. The CCT of eukaryotic cytosol is composed of eight different subunit species, TCP-1 α , β , γ , δ , ϵ , ζ , η and θ , each encoded by a different gene. Two ζ subunits have been described: TCP-1 ζ (also designated TCP-1 ξ 1) and TCP-1 ζ 2. TCP-1 subunits are proposed to have independent functions in folding its *in vivo* substrates, the actins and tubulins. TCP-1 was first identified in the mouse as relevant for tail-less and embryonic lethal phenotypes. Sequences homologous to TCP-1 have been isolated in several other species, and the yeast TCP-1 has been shown to encode a molecular chaperone for Actin and Tubulin. TCP-1 found in mammalian cells and yeast plays an important role in the folding of cytosolic proteins.

REFERENCES

1. Ahnert, V., et al. 1996. Cucumber t complex protein. Molecular cloning, bacterial expression and characterization within a 22-S cytosolic complex in cotyledons and hypocotyls. *Eur. J. Biochem.* 235: 114-119.
2. Iijima, M., et al. 1998. A *Dictyostelium discoideum* homologue to TCP-1 is essential for growth and development. *Gene* 213: 101-106.
3. Ritco-Vonsovici, M. and Willison, K.R. 2000. Defining the eukaryotic cytosolic chaperonin-binding sites in human tubulins. *J. Mol. Biol.* 304: 81-98.
4. Hynes, G.M. and Willison, K.R. 2000. Individual subunits of the eukaryotic cytosolic chaperonin mediate interactions with binding sites located on subdomains of β -Actin. *J. Biol. Chem.* 275: 18985-18994.
5. Campos, E.G. and Hamdan, F.F. 2000. Cloning of the chaperonin t complex polypeptide 1 gene from *Schistosoma mansoni* and studies of its expression levels under heat shock and oxidative stress. *Parasitol. Res.* 86: 253-258.
6. Yokota, S.I., et al. 2000. Upregulation of cytosolic chaperonin CCT subunits during recovery from chemical stress that causes accumulation of unfolded proteins. *Eur. J. Biochem.* 267: 1658-1664.

CHROMOSOMAL LOCATION

Genetic locus: CCT7 (human) mapping to 2p13.2; Cct7 (mouse) mapping to 6 C3.

SOURCE

TCP-1 η (C-6) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 474-512 near the C-terminus of TCP-1 η of human origin.

PRODUCT

Each vial contains 200 μ g IgG_{2b} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-271439 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

APPLICATIONS

TCP-1 η (C-6) is recommended for detection of TCP-1 η of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

TCP-1 η (C-6) is also recommended for detection of TCP-1 η in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for TCP-1 η siRNA (h): sc-43449, TCP-1 η siRNA (m): sc-43450, TCP-1 η shRNA Plasmid (h): sc-43449-SH, TCP-1 η shRNA Plasmid (m): sc-43450-SH, TCP-1 η shRNA (h) Lentiviral Particles: sc-43449-V and TCP-1 η shRNA (m) Lentiviral Particles: sc-43450-V.

Molecular Weight of TCP-1 η : 58 kDa.


Positive Controls: F9 cell lysate: sc-2245, K-562 whole cell lysate: sc-2203 or HL-60 whole cell lysate: sc-2209.

RECOMMENDED SUPPORT REAGENTS


To ensure optimal results, the following support reagents are recommended:

- 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.
- 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).
- 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



TCP-1 η (C-6): sc-271439. Western blot analysis of TCP-1 η expression in HL-60 (A) and K-562 (B) whole cell lysates.



TCP-1 η (C-6): sc-271439. Immunofluorescence staining of methanol-fixed HeLa cells showing cytoplasmic localization.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.