

FGF-10 (A-15): sc-27147

BACKGROUND

Fibroblast growth factor-1 (FGF-1), also designated acidic FGF, and fibroblast growth factor-2 (FGF-2), also designated basic FGF, are members of a family of growth factors that stimulate proliferation of cells of mesenchymal, epithelial and neuroectodermal origin. Additional members of the FGF family include the oncogenes FGF-3 (Int2) and FGF-4 (hst/Kaposi), FGF-5, FGF-6, FGF-7 (KGF), FGF-8 (AIGF), FGF-9 (GAF) and FGF-10–FGF-23. Members of the FGF family share 30-55% amino acid sequence identity and similar gene structure, and are capable of transforming cultured cells when overexpressed in transfected cells. Cellular receptors for FGFs are members of a second multigene family including four tyrosine kinases, designated Flg (FGFR-1), Bek (FGFR-L), TKF and FGFR-3.

REFERENCES

- Moore, R., Casey, G., Brookes, S., Dixon, M., Peters, G. and Dickson, C. 1986. Sequence, topography and protein coding potential of mouse int-2: a putative oncogene activated by mouse mammary tumor virus. *EMBO J.* 5: 919-924.
- Delli Bovi, P., Curatola, A.M., Kern, F.G., Greco, A., Ittmann, M. and Basilico, C. 1987. An oncogene isolated by transfection of Kaposi's sarcoma DNA encodes a growth factor that is a member of the FGF family. *Cell* 50: 729-737.
- Zhan, X., Bates, B., Hu, X. and Goldfarb, M. 1988. The human FGF-5 oncogene encodes a novel protein related to fibroblast growth factors. *Mol. Cell Biol.* 8: 3487-3495.
- Marics, I., Adelaide, J., Raybaud, F., Mattei, M., Coulier, F., Planche, J., Lapeyriere, O. and Birnbaum, D. 1989. Characterization of the HST-related FGF.6 gene, a new member of the fibroblast growth factor gene family. *Oncogene* 4: 335-340.
- Rifkin, D.B. and Moscatelli, D. 1989. Recent developments in the cell biology of fibroblast growth factor. *J. Cell Biol.* 109: 1-6.
- Tanaka, A., Miyamoto, K., Minamino, N., Takeda, M., Sato, B., Matsuo, H. and Matsumoto, K. 1992. Cloning and characterization of an androgen-induced growth factor essential for the androgen-dependent growth of mouse mammary carcinoma cells. *Proc. Natl. Acad. Sci. USA* 89: 8928-8932.
- Miyamoto, M., Naruo, K., Seko, C., Matsumoto, S., Kondo, T. and Kurokawa, T. 1993. Molecular cloning of a novel cytokine cDNA encoding the ninth member of the fibroblast growth factor family, which has a unique secretion property. *Mol. Cell Biol.* 13: 4251-4259.
- Beer, H.D., Florence, C., Dammeier, J., McGuire, L., Werner, S. and Duan, D.R. 1997. Mouse fibroblast growth factor 10: cDNA cloning, protein characterization, and regulation of mRNA expression. *Oncogene* 15: 2211-2218.

CHROMOSOMAL LOCATION

Genetic locus: FGF10 (human) mapping to 5p12; Fgf10 (mouse) mapping to 13 D2.3.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

SOURCE

FGF-10 (A-15) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of FGF-10 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-27147 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

FGF-10 (A-15) is recommended for detection of precursor and mature FGF-10 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

FGF-10 (A-15) is also recommended for detection of precursor and mature FGF-10 in additional species, including equine and porcine.

Suitable for use as control antibody for FGF-10 siRNA (h): sc-39462, FGF-10 siRNA (m): sc-39463, FGF-10 shRNA Plasmid (h): sc-39462-SH, FGF-10 shRNA Plasmid (m): sc-39463-SH, FGF-10 shRNA (h) Lentiviral Particles: sc-39462-V and FGF-10 shRNA (m) Lentiviral Particles: sc-39463-V.

Molecular Weight of FGF-10: 19 kDa.

Positive Controls: human malignant glioma tissue extract.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.



MONOS
Satisfaction
Guaranteed

Try **FGF-10 (3C7): sc-293208**, our highly recommended monoclonal alternative to FGF-10 (A-15).