



Ku70 (aN-20): sc-27167

BACKGROUND

In higher organisms such as mammals and plants, DNA double-strand breaks are repaired preferentially by non-homologous end joining (NHEJ) rather than by homologous recombination. The NHEJ pathway is mediated by Ku, a heterodimer of approximately 70 and 80 kDa subunits, which contributes to various aspects of the metabolism of DNA ends in eukaryotic cells. This heterodimer binds to double-stranded telomeric and non-telomeric DNA sequences. The *Arabidopsis* Ku heterodimer also possesses single-stranded DNA-dependent ATPase and ATP-dependent DNA helicase activities. Ku80 can translocate to the nucleus using its own NLS independent of the translocation of Ku70.

REFERENCES

- Riha, K., et al. 2002. Telomere length deregulation and enhanced sensitivity to genotoxic stress in *Arabidopsis* mutants deficient in Ku70. *Embo. J.* 21: 2819-2826.
- Bundock, P., et al. 2002. Increased telomere length and hypersensitivity to DNA damaging agents in an *Arabidopsis* KU70 mutant. *Nucleic. Acids. Res.* 30: 3395-3400.
- Tamura, K., et al. 2002. Identification of Ku70 and Ku80 homologues in *Arabidopsis thaliana*: evidence for a role in the repair of DNA double-strand breaks. *Plant J.* 29: 771-781.
- West, C.E., et al. 2002. Disruption of the *Arabidopsis* AtKu80 gene demonstrates an essential role for AtKu80 protein in efficient repair of DNA double-strand breaks *in vivo*. *Plant J.* 31: 517-528.
- Riha, K., et al. 2003. Ku is required for telomeric C-rich strand maintenance but not for end-to-end chromosome fusions in *Arabidopsis*. *Proc. Natl. Acad. Sci. USA.* 100: 611-615.
- Gallego, M.E., et al. 2003. Telomerase dependence of telomere lengthening in Ku80 mutant *Arabidopsis*. *Plant Cell.* 15: 782-789.
- Gallego, M.E., et al. 2003. Ku80 plays a role in non-homologous recombination but is not required for T-DNA integration in *Arabidopsis*. *Plant J.* 35: 557-565.
- Heacock, M., et al. 2004. Molecular analysis of telomere fusions in *Arabidopsis*. Multiple pathways for chromosome end-joining. *Embo. J.* 23: 2304-2313.
- Puizina, J., et al. 2004. Mre11 deficiency in *Arabidopsis* is associated with chromosomal instability in somatic cells and Spo11-dependent genome fragmentation during meiosis. *Plant Cell.* 16: 1968-1978.
- Kuchar, M., et al. 2004. Interactions of putative telomere-binding proteins in *Arabidopsis thaliana*: identification of functional TRF2 homolog in plants. *FEBS. Letts.* 578: 311-315.

SOURCE

Ku70 (aN-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the N-terminus of Ku70 of *Arabidopsis thaliana* origin.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-27167 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

Ku70 (aN-20) is recommended for detection of Ku70 of *Arabidopsis thaliana* origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Molecular Weight of Ku70: 70 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.