

GS1 (H-2): sc-271692

BACKGROUND

The Adiponutrin family consists of Adiponutrin (ADPN), adipocyte triglyceride lipase (ATGL, also designated Desnutrin), GS1, GS2, GS2-like and PNPLA1. ADPN, ATGL and GS2 display lipase activity, which is dependent upon the presence of an activated serine residue. GS1, also designated DXF68S1E or haloacid dehalogenase-like hydrolase domain containing 1A (HDHD1A), is a 214-amino acid protein that is detected in human placenta and fibroblasts. The gene which encodes for GS1, HDHD1A, is of interest because it is an X-linked gene that escapes X-inactivation. This characteristic of the HDHD1A gene is particularly important in the understanding of human X chromosome structural organization as well as the mechanism of X-inactivation.

REFERENCES

- Salido, E.C., et al. 1992. The human enamel protein gene amelogenin is expressed from both the X and the Y chromosomes. *Am. J. Hum. Genet.* 50: 303-316.
- Yen, P.H., et al. 1993. Isolation of a new gene from the distal short arm of the human X chromosome that escapes X-inactivation. *Hum. Mol. Genet.* 1: 47-52.
- Soehnge, H., et al. 1997. Cloning and sequencing of Ribosomal Protein L27a and a gene similar to human GS1 in *Drosophila*. *Gene* 185: 257-263.
- Online Mendelian Inheritance in Man, OMIM™. 1998. Johns Hopkins University, Baltimore, MD. MIM Number: 306480. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- van Noort, V., et al. 2003. Predicting gene function by conserved co-expression. *Trends Genet.* 19: 238-242.
- Wieland, I., et al. 2003. Further delineation of Wittwer syndrome and refinement of the mapping region. *Am. J. Med. Genet. A* 116A: 57-60.
- Gerhard, D.S., et al. 2004. The status, quality, and expansion of the NIH full-length cDNA project: the mammalian gene collection (MGC). *Genome Res.* 14: 2121-2127.

CHROMOSOMAL LOCATION

Genetic locus: PUDP (human) mapping to Xp22.31; Hdhd1a (mouse) mapping to 18 D1.

SOURCE

GS1 (H-2) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 207-234 at the C-terminus of GS1 of mouse origin.

PRODUCT

Each vial contains 200 µg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-271692 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

APPLICATIONS

GS1 (H-2) is recommended for detection of GS1 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for GS1 siRNA (h): sc-60768, GS1 siRNA (m): sc-60769, GS1 shRNA Plasmid (h): sc-60768-SH, GS1 shRNA Plasmid (m): sc-60769-SH, GS1 shRNA (h) Lentiviral Particles: sc-60768-V and GS1 shRNA (m) Lentiviral Particles: sc-60769-V.

Molecular Weight of GS1: 24 kDa.


Positive Controls: HeLa whole cell lysate: sc-2200, K-562 whole cell lysate: sc-2203 or COLO 205 whole cell lysate: sc-364177.

RECOMMENDED SUPPORT REAGENTS


To ensure optimal results, the following support reagents are recommended:

- 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.
- 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).
- 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.
- 4) Immunohistochemistry: use m-IgGκ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

DATA



GS1 (H-2): sc-271692. Western blot analysis of GS1 expression in HeLa (A), K-562 (B), COLO 205 (C), SW480 (D) and NIH/3T3 (E) whole cell lysates.



GS1 (H-2): sc-271692. Immunoperoxidase staining of formalin fixed, paraffin-embedded human placenta tissue showing cytoplasmic staining of trophoblastic cells.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.