

NTF2 (B-7): sc-271693

BACKGROUND

Protein transport across the nucleus is a selective, multistep process involving several cytoplasmic factors. Proteins must be recognized as import substrates, dock at the nuclear pore complex and translocate across the nuclear envelope in an ATP-dependent fashion. Two cytosolic factors centrally involved in the recognition and docking process are the karyopherin- α and karyopherin- β proteins. The karyopherin holoenzyme is a heterodimer of α and β subunits. Karyopherin- α functions in the recognition and targeting of substrates destined for nuclear import, while karyopherin- β serves as an adapter, tethering the karyopherin- α /substrate complex to docking proteins on the nuclear envelope termed nucleoporins. p62 glycoprotein is one such nucleoporin, and is not only involved in the nuclear import of proteins, but also the export of nascent mRNA strands. An additional protein, NTF2 (nuclear transport factor 2), interacts with nucleoporin p62 as a homodimer, and may be an obligate component of functional p62.

REFERENCES

1. Buss, F., et al. 1995. Macromolecular interactions in the nucleoporin p62 complex of rat nuclear pores: binding of nucleoporin p54 to the rod domain of p62. *J. Cell Biol.* 128: 251-261.
2. Paschal, B.M., et al. 1995. Identification of NTF2, a cytosolic factor for nuclear import that interacts with nuclear pore complex protein p62. *J. Cell Biol.* 129: 925-937.
3. Moroianu, J., et al. 1995. Previously identified protein of uncertain function is karyopherin α and together with karyopherin β docks import substrate at nuclear pore complexes. *Proc. Natl. Acad. Sci. USA* 92: 2008-2011.

CHROMOSOMAL LOCATION

Genetic locus: NUTF2 (human) mapping to 16q22.1; Nutf2 (mouse) mapping to 8 D3.

SOURCE

NTF2 (B-7) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 97-127 at the C-terminus of nuclear transport factor 2 (NTF2) of human origin.

PRODUCT

Each vial contains 200 μ g IgG_{2b} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

NTF2 (B-7) is available conjugated to agarose (sc-271693 AC), 500 μ g/0.25 ml agarose in 1 ml, for IP; to HRP (sc-271693 HRP), 200 μ g/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-271693 PE), fluorescein (sc-271693 FITC), Alexa Fluor® 488 (sc-271693 AF488), Alexa Fluor® 546 (sc-271693 AF546), Alexa Fluor® 594 (sc-271693 AF594) or Alexa Fluor® 647 (sc-271693 AF647), 200 μ g/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-271693 AF680) or Alexa Fluor® 790 (sc-271693 AF790), 200 μ g/ml, for Near-Infrared (NIR) WB, IF and FCM.

Blocking peptide available for competition studies, sc-271693 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

APPLICATIONS

NTF2 (B-7) is recommended for detection of NTF2 p10 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for NTF2 siRNA (h): sc-36105, NTF2 siRNA (m): sc-36106, NTF2 shRNA Plasmid (h): sc-36105-SH, NTF2 shRNA Plasmid (m): sc-36106-SH, NTF2 shRNA (h) Lentiviral Particles: sc-36105-V and NTF2 shRNA (m) Lentiviral Particles: sc-36106-V.


Molecular Weight of NTF2: 14 kDa.

Positive Controls: COLO 320DM cell lysate: sc-2226, HeLa whole cell lysate: sc-2200 or K-562 whole cell lysate: sc-2203.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



NTF2 (B-7): sc-271693. Western blot analysis of NTF2 expression in HeLa (A) and COLO 320DM (B) whole cell lysates.



NTF2 (B-7): sc-271693. Immunofluorescence staining of methanol-fixed HeLa cells showing cytoplasmic localization.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA