

KAT I (B-8): sc-271709

BACKGROUND

Kynurenine aminotransferases KAT I, KAT II and KAT III belong to the class-I pyridoxal-phosphate-dependent aminotransferase family. KAT I is a cytoplasmic protein involved in glutamine catabolism. KAT I functions in the catalysis of the transamination of L-kinurenine to form kynurenic acid, a neuroprotective and anticonvulsant metabolite of tryptophan. Kynurenic acid is involved in synaptic transmission and has been implicated in a number of neurological disorders including schizophrenia and Huntington's disease. KAT I also functions in the metabolism of cysteine conjugates in some halogenated alkenes and alkanes to form reactive metabolites. KAT I has three isoforms. Isoform 1 is the full length protein, isoform 2 lacks amino acids 68-117 and isoform 3 lacks amino acids 251-422. Based on sequence similarity, KAT I is thought to function as a homodimer.

REFERENCES

1. Baran, H., et al. 1996. Increased kynurenic acid levels and decreased brain kynurenine aminotransferase I in patients with Down syndrome. *Life Sci.* 58: 1891-1899.
2. Tamburin, M., et al. 1999. Kynurenine aminotransferase I (KAT I) isoform gene expression in the rat brain: an *in situ* hybridization study. *Neuroreport* 10: 61-65.
3. Milart, P., et al. 2001. Kynurenine aminotransferase I activity in human placenta. *Placenta* 22: 259-261.
4. Kwok, J.B., et al. 2002. A missense mutation in kynurenine aminotransferase-1 in spontaneously hypertensive rats. *J. Biol. Chem.* 277: 35779-35782.
5. Rejda, R., et al. 2003. Ontogenic changes of kynurenine aminotransferase I activity and its expression in the chicken retina. *Vision Res.* 43: 1513-1517.
6. Knyihár-Csillik, E., et al. 2004. Decreased expression of kynurenine aminotransferase-I (KAT I) in the substantia nigra of mice after 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP) treatment. *Neuroscience* 126: 899-914.
7. Rossi, F., et al. 2004. Crystal structure of human kynurenine aminotransferase I. *J. Biol. Chem.* 279: 50214-50220.

CHROMOSOMAL LOCATION

Genetic locus: CCBL1 (human) mapping to 9q34.11; Ccbl1 (mouse) mapping to 2 B.

SOURCE

KAT I (B-8) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 341-377 within an internal region of KAT I of human origin.

PRODUCT

Each vial contains 200 µg IgG_{2a} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-271709 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

APPLICATIONS

KAT I (B-8) is recommended for detection of KAT I isoforms 1 and 2 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for KAT I siRNA (h): sc-105587, KAT I siRNA (m): sc-77396, KAT I shRNA Plasmid (h): sc-105587-SH, KAT I shRNA Plasmid (m): sc-77396-SH, KAT I shRNA (h) Lentiviral Particles: sc-105587-V and KAT I shRNA (m) Lentiviral Particles: sc-77396-V.


Molecular Weight of KAT I: 48 kDa.

Positive Controls: KAT I (m): 293T Lysate: sc-127032 or HeLa whole cell lysate: sc-2200.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgGκ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

DATA



KAT I (B-8): sc-271709. Western blot analysis of KAT I expression in non-transfected: sc-117752 (A) and mouse KAT I transfected: sc-127032 (B) 293T whole cell lysates.



KAT I (B-8): sc-271709. Immunoperoxidase staining of formalin fixed, paraffin-embedded human skeletal muscle tissue showing cytoplasmic staining of myocytes.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.