

RFX5 (C-3): sc-271756

BACKGROUND

EP is a regulatory enhancer element found in several promoters on viral genes, and similar sites are also present in cellular genes, including the MIF-1 binding site (MIE) of the human c-Myc gene, the X box of MHC class II promoters and a binding site in the proliferating cell nuclear antigen promoter. The EP sites present in the X box of MHC class II promoters are distinctly nonpalindromic sequences that contain only a single EP-homologous half-site. The EP element is bound by an ubiquitous nuclear protein complex that consists of homo- and heterodimers involving the RFX1, RFX2 and RFX3 proteins. The RFX proteins represent an essential class II transcription factor family that shares several conserved regions, including the centrally located DNA-binding domain (DBD) and the D region found in the C-terminal part of these proteins which facilitates dimerization. RFX complexes can activate the enhancer elements of several HBV genes and also promote the induction of MHC class II genes in response to interferon- γ stimulation. Two additional subunits, RFX5, RFX-B/Ank, are also involved in the RFX complexes, yet they bind additional elements in the X1 half of the X box.

REFERENCES

1. Dikstein, R., et al. 1990. Functional organization of the hepatitis B virus enhancer. *Mol. Cell. Biol.* 10: 3682-3689.
2. Fontes, J.D., et al. 1997. Assembly of functional regulatory complexes on MHC class II promoters *in vivo*. *J. Mol. Biol.* 270: 336-345.
3. Katan, Y., et al. 1997. The transcriptional activation and repression domains of RFX1, a context-dependent regulator, can mutually neutralize their activities. *Nucleic Acids Res.* 25: 3621-3628.

CHROMOSOMAL LOCATION

Genetic locus: RFX5 (human) mapping to 1q21.3.

SOURCE

RFX5 (C-3) is a mouse monoclonal antibody raised against amino acids 161-460 mapping within an internal region of RFX5 of human origin.

PRODUCT

Each vial contains 200 μ g IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin. Also available as TransCruz reagent for Gel Supershift and ChIP applications, sc-271756 X, 200 μ g/0.1 ml.

RFX5 (C-3) is available conjugated to agarose (sc-271756 AC), 500 μ g/0.25 ml agarose in 1 ml, for IP; to HRP (sc-271756 HRP), 200 μ g/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-271756 PE), fluorescein (sc-271756 FITC), Alexa Fluor[®] 488 (sc-271756 AF488), Alexa Fluor[®] 546 (sc-271756 AF546), Alexa Fluor[®] 594 (sc-271756 AF594) or Alexa Fluor[®] 647 (sc-271756 AF647), 200 μ g/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor[®] 680 (sc-271756 AF680) or Alexa Fluor[®] 790 (sc-271756 AF790), 200 μ g/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor[®] is a trademark of Molecular Probes, Inc., Oregon, USA

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

RFX5 (C-3) is recommended for detection of RFX5 of human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for RFX5 siRNA (h): sc-37747, RFX5 shRNA Plasmid (h): sc-37747-SH and RFX5 shRNA (h) Lentiviral Particles: sc-37747-V.

RFX5 (C-3) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.


Molecular Weight of RFX5: 75 kDa.

Positive Controls: Jurkat nuclear extract: sc-2132, Ramos nuclear extract: sc-2153 or Raji whole cell lysate: sc-364236.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

DATA



RFX5 (C-3): sc-271756. Western blot analysis of RFX5 expression in Jurkat (A) and Ramos (B) nuclear extracts and U-698-M whole cell lysate (C).

RFX5 (C-3): sc-271756. Western blot analysis of RFX5 expression in Jurkat nuclear extract (A) and Raji whole cell lysate (B).

SELECT PRODUCT CITATIONS

1. Prusty, B.K., et al. 2018. Peptidase inhibitor 15 (PI15) regulates chlamydial CPAF activity. *Front. Cell. Infect. Microbiol.* 8: 183.
2. Chirichella, M., et al. 2022. RFX transcription factors control a miR-150/PDAP1 axis that restrains the proliferation of human T cells. *PLoS Biol.* 20: e3001538.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.