SANTA CRUZ BIOTECHNOLOGY, INC.

ARK5 (A-9): sc-271827



BACKGROUND

5'-AMP-activated protein kinase, known as AMPK, is a heterotrimeric complex that protects cells from stresses that cause ATP depletion by switching off ATP-consuming biosynthetic pathways. Activated AMPK phosphorylates and regulates hydroxymethylglutaryl-CoA reductase and acetyl-CoA carboxylase, which are key regulatory enzymes of sterol synthesis and fatty acid synthesis, respectively. The AMPK-related kinase 5 (ARK5) is a tumor progression-associated factor that is directly phosphorylated by Akt at Ser 600, which is in its regulatory domain. NDR2, a serine/threonine protein kinase, phosphorylates and activates ARK5 during Insulin-like growth factor (IGF)-I signaling in order to promote cell survival. This activation may be involved in the invasion of colorectal cancer cell lines. Furthermore, ARK5 is a transcriptional target of the Large-MAF family and may have a role in mediating the aggressiveness of c-Maf- and MafB-expressing myelomas.

REFERENCES

- Suzuki, A., et al. 2003. ARK5 suppresses the cell death induced by nutrient starvation and death receptors via inhibition of caspase 8 activation, but not by chemotherapeutic agents or UV irradiation. Oncogene 22: 6177-6182.
- Suzuki, A., et al. 2004. ARK5 is a tumor invasion-associated factor downstream of Akt signaling. Mol. Cell. Biol. 24: 3526-3535.
- Suzuki, A., et al. 2004. Regulation of caspase-6 and FLIP by the AMPK family member ARK5. Oncogene 23: 7067-7075.
- Fisher, J.S., et al. 2005. Muscle contractions, AICAR, and Insulin cause phosphorylation of an AMPK-related kinase. Am. J. Physiol. Endocrinol. Metab. 289: E986-E992.

CHROMOSOMAL LOCATION

Genetic locus: NUAK1 (human) mapping to 12q23.3.

SOURCE

ARK5 (A-9) is a mouse monoclonal antibody raised against amino acids 526-661 mapping at the C-terminus of ARK5 of human origin.

PRODUCT

Each vial contains 200 μg IgG_1 kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

ARK5 (A-9) is available conjugated to agarose (sc-271827 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-271827 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-271827 PE), fluorescein (sc-271827 FITC), Alexa Fluor[®] 488 (sc-271827 AF488), Alexa Fluor[®] 546 (sc-271827 AF546), Alexa Fluor[®] 594 (sc-271827 AF594) or Alexa Fluor[®] 647 (sc-271827 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor[®] 680 (sc-271827 AF680) or Alexa Fluor[®] 790 (sc-271827 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

ARK5 (A-9) is recommended for detection of ARK5 of human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for ARK5 siRNA (h): sc-60203, ARK5 shRNA Plasmid (h): sc-60203-SH and ARK5 shRNA (h) Lentiviral Particles: sc-60203-V.


Molecular Weight of ARK5: 74 kDa.


Positive Controls: ARK5 (h): 293T Lysate: sc-173768, Saos-2 whole cell lysate: sc-2235 or A549 whole cell lysate: sc-2413.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

DATA





ARK5 (A-9): sc-271827. Western blot analysis of ARK5 expression in Hep G2 $({\rm A}),$ Saos-2 $({\rm B})$ and A549 $({\rm C})$ whole cell lysates.

ARK5 (A-9): sc-271827. Western blot analysis of ARK5 expression in non-transfected: sc-117752 (**A**) and human ARK5 transfected: sc-173768 (**B**) 293T whole cell lysates.

SELECT PRODUCT CITATIONS

 Chen, M., et al. 2019. LINC00958 promotes the malignancy of nasopharyngeal carcinoma by sponging microRNA-625 and thus upregulating NUAK1. Onco Targets Ther. 12: 9277-9290.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA