



The Power to Question

Mesothelin (V-16): sc-27714

BACKGROUND

Mesothelin is a glycosylphosphatidylinositol-linked cell-surface molecule expressed in the mesothelial lining of the body cavities and in many tumor cells. Mesothelin is a tumor antigen on the surface of human ovarian cancers and mesotheliomas. Mesothelin immunoreactivity is high in cancers of the ovary (serous papillary, endometrioid and undifferentiated) and pancreas, with less frequent staining seen in adenocarcinomas of the endometrium, lung and stomach/esophagus. In adult mouse tissues the Mesothelin transcript is present in lung, heart, spleen, liver, kidney and testis.

REFERENCES

- Chang, K., et al. 1996. Molecular cloning of Mesothelin, a differentiation antigen present on mesothelium, mesotheliomas, and ovarian cancers. Proc. Natl. Acad. Sci. USA 93: 136-140.
- Chowdhury, P.S., et al. 1997. Isolation of anti-Mesothelin antibodies from a phage display library. Mol. Immunol. 34: 9-20.
- Hassan, R., et al. 1999. 111Indium-labeled monoclonal antibody K1: biodistribution study in nude mice bearing a human carcinoma xenograft expressing mesothelin. Int. J. Cancer. 80: 559-563.
- Bera, T.K., et al. 2000. Mesothelin is not required for normal mouse development or reproduction. Mol. Cell. Biol. 20: 2902-2906.
- Frierson, H.F., Jr., et al. 2003. Large-scale molecular and tissue microarray analysis of Mesothelin expression in common human carcinomas. Hum. Pathol. 34: 605-609.
- Rump, A., et al. 2004. Binding of ovarian cancer antigen CA125/MUC16 to Mesothelin mediates cell adhesion. J. Biol. Chem. 279: 9190-9198.

CHROMOSOMAL LOCATION

Genetic locus: MSLN (human) mapping to 16p13.3.

SOURCE

Mesothelin (V-16) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of Mesothelin of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-27714 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

APPLICATIONS

Mesothelin (V-16) is recommended for detection of precursor and mature Mesothelin of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for Mesothelin siRNA (h): sc-45386, Mesothelin shRNA Plasmid (h): sc-45386-SH and Mesothelin shRNA (h) Lentiviral Particles: sc-45386-V.

Molecular Weight of Mesothelin precursor: 69 kDa.


Molecular Weight of mature Mesothelin: 40 kDa.

Positive Controls: AN3 CA cell lysate: sc-24662, ES-2 cell lysate: sc-24674 or Ca Ski whole cell lysate.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



Mesothelin (V-16): sc-27714. Western blot analysis of Mesothelin expression in ES-2 (**A**) and Ca Ski (**B**) whole cell lysates.

Mesothelin (V-16): sc-27714. Western blot analysis of Mesothelin expression in AN3 CA whole cell lysate.

SELECT PRODUCT CITATIONS

- Johnson, M.D., et al. 2008. Mesothelin expression in the leptomeninges and meningiomas. J. Histochem. Cytochem. 56: 579-585.

RESEARCH USE

For research use only, not for use in diagnostic procedures.