Pik1 (yN-15): sc-27738



The Power to Question

BACKGROUND

Phosphatidylinositol 4-kinases are molecules that function as important second messengers in various cellular processes. Pik1 is a 1066 amino acid phosphatidylinositol 4-kinase present in yeast that localizes to the cytoplasmic puncta and the nucleus. Pik1 catalyzes the first committed step in the pathway that produces the second messenger inositol-1,4,5-trisphosphate, which is used during signal transduction. Pik1 is necessary for normal cellular secretion, Golgi and vacuole membrane dynamics, as well as endocytosis. Pik1 may also control cytokinesis via the Actin cytoskeleton. Frequenin (Frq1), a 190-residue N-myristoylated calcium-binding protein, associates tightly with the N terminus of Pik1, and both are essential for viability and growth.

REFERENCES

- 1. Flanagan, C.A., Schnieders, E.A., Emerick, A.W., Kunisawa, R., Admon, A. and Thorner, J. 1993. Phosphatidylinositol 4-kinase: gene structure and requirement for yeast cell viability. Science 262: 1444-1448.
- Garcia-Bustos, J.F., Marini, F., Stevenson, I., Frei, C. and Hall, M.N. 1994.
 Pik1, an essential phosphatidylinositol 4-kinase associated with the yeast nucleus. EMBO J. 13: 2352-2361.
- Hama, H., Schnieders, E.A., Thorner, J., Takemoto, J.Y. and DeWald, D.B. 1999. Direct involvement of phosphatidylinositol 4-phosphate in secretion in the yeast *Saccharomyces cerevisiae*. J. Biol. Chem. 274: 34294-34300.
- Audhya, A., Foti, M. and Emr, S.D. 2000. Distinct roles for the yeast phosphatidy in secretion, cell growth, and organelle membrane dynamics. Mol. Biol. Cell 11: 2673-2689.
- Walch-Solimena, C. and Novick, P. 2000. The yeast phosphatidylinositol-4-OH kinase Pik1 regulates secretion at the Golgi. Nat. Cell Biol. 1: 523-525.
- Huttner, I.G., Strahl, T., Osawa, M., King, D.S., Ames, J.B. and Thorner, J. 2003. Molecular interactions of yeast frequenin (Frq1) with the phosphatidylinositol 4-kinase isoform, Pik1. J. Biol. Chem. 278: 4862-4874.
- Strahl, T., Grafelmann, B., Dannenberg, J., Thorner, J. and Pongs, O. 2003. Conservation of regulatory function in calcium-binding proteins: human frequenin (neuronal calcium sensor-1) associates productively with yeast phosphatidylinositol 4-kinase isoform, Pik1. J. Biol. Chem. 278: 49589-49599.
- Strahl, T., Hama, H., DeWald, D.B. and Thorner, J. 2005. Yeast phosphatidylinositol 4-kinase, Pik1, has essential roles at the Golgi and in the nucleus. J. Cell Biol. 171: 967-979.
- 9. Minogue, S., Waugh, M.G., De Matteis, M.A., Stephens, D.J., Berditchevski, F. and Hsuan, J.J. 2006. Phosphatidylinositol 4-kinase is required for endosomal trafficking and degradation of the EGF receptor. J. Cell Sci.119 (Pt. 3): 571-581.

SOURCE

Pik1 (yN-15) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of Pik1 of *Saccharomyces cerevisiae* origin.

PRODUCT

Each vial contains 200 μg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-27738 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

Pik1 (yN-15) is recommended for detection of Pik1 of *Saccharomyces cerevisiae* origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

Santa Cruz Biotechnology, Inc. 1.800.457.3801 831.457.3801 **Europe** +00800 4573 8000 49 6221 4503 0 **www.scbt.com**