SEMA3C (V-20): sc-27797



The Power to Question

BACKGROUND

SEMA3C (also designated Semaphorin 3C, Semaphorin E, SEMAE, SemE, sema domain, immunoglobulin domain (Ig) and short basic domain, secreted) is an ubiquitous protein of approximately 70 kDa that mediates axonal guidance, repulsive gradients, induction of growth cone collapse and cell survival/ death. Secreted SEMA3C favors survival and neuritogenesis of cultured cerebellar granule neurons (CGNs). SEMA3C from macrophages and fibroblasts that selectively directs against sympathetic nerve fibers may be one element responsible for reduced sympathetic innervation in rheumatoid arthritis tissue. SEMA3C mutant mice die within hours after birth from congenital cardiovascular defects consisting of interruption of the aortic arch and improper septation of the cardiac outflow tract. SEMA3C is expressed in the cardiac outflow tract as neural crest cells and promotes crest cell migration into the proximal cardiac outflow tract. Semaphorins constitute a family of molecules sharing a common extracellular domain (Semaphorin domain). The family includes several types of secreted and membrane-associated molecules that are grouped into eight subclasses (subclasses 1-7 and viral Semaphorins).

REFERENCES

- Kolodkin, A.L., et al. 1993. The Semaphorin genes encode a family of transmembrane and secreted growth cone guidance molecules. Cell 75: 1389-1399.
- Puschel, A.W., et al. 1995. Murine Semaphorin D/Collapsin is a member of a diverse gene family and creates domains inhibitory for axonal extension. Neuron 14: 941-948.
- 3. Dodd, J., et al. 1995. Axon guidance: a compelling case for repelling growth cones. Cell 81: 471-474.
- 4. Matthes, D.J., et al. 1995. Semaphorin II can function as a selective inhibitor of specific synaptic arborizations. Cell 81: 631-639.
- Messersmith, E.K., et al. 1995. Semaphorin III can function as a selective chemorepellent to pattern sensory projections in the spinal cord. Neuron 14: 949-959.
- Wright, D.E., et al. 1995. The guidance molecule Semaphorin III is expressed in regions of spinal cord and periphery avoided by growing sensory axons. J. Comp. Neurol. 361: 321-333.

CHROMOSOMAL LOCATION

Genetic locus: SEMA3C (human) mapping to 7q21-q31; Sema3c (mouse) mapping to 5 A3.

SOURCE

SEMA3C (V-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of SEMA3C of human origin.

PRODUCT

Each vial contains 200 μg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-27797 P, (100 μg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

SEMA3C (V-20) is recommended for detection of precursor and mature SEMA3C of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for SEMA3C siRNA (h): sc-44091 and SEMA3C siRNA (m): sc-44381.

Molecular Weight of SEMA3C: 70 kDa.

Positive Controls: rat brain extract: sc-2392.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

Santa Cruz Biotechnology, Inc. 1.800.457.3801 831.457.3800 fax 831.457.3801 **Europe** +00800 4573 8000 49 6221 4503 0 **www.scbt.com**