

SMC1 β (C-16): sc-27961

BACKGROUND

The SMC (structural maintenance of chromosomes) family of proteins form heterodimeric complexes that modulate sister chromatid cohesion and chromosome condensation for mitosis. The two distinct classes of SMC protein complexes are comprised of SMC1 (also designated SB1.8) with SMC3 (also designated HCAP for human chromosome-associated protein and Bamacan for the secreted proteoglycan), and SMC2 (also designated hCAP-E) with SMC4 (also designated hCAP-C). The SMC1/SMC3 complex is required for metaphase progression in mitotic cells and functions independently of the SMC2/SMC4 complex during the cell cycle. SMC1 is ubiquitously expressed in various human tissues, including thymus, testis, and colon. SMC3 is expressed as a nuclear protein in the colon, but can also occur as a secreted proteoglycan expressed in testis and brain. The secreted proteoglycan contains several glycosylation sites and is thought to play a role in basement membrane physiology.

REFERENCES

1. Strunnikov, A.V., et al. 1993. SMC1: an essential yeast gene encoding a putative head-rod-tail protein is required for nuclear division and defines a new ubiquitous protein family. *J. Cell Biol.* 123: 1635-1648.
2. Rocques, P.J., et al. 1995. The human SB1.8 gene (DXS423E) encodes a putative chromosome segregation protein conserved in lower eukaryotes and prokaryotes. *Hum. Mol. Genet.* 4: 243-249.
3. Ljubimov, A.V., et al. 1996. Basement membrane abnormalities in human eyes with diabetic retinopathy. *J. Histochem. Cytochem.* 44: 1469-1479.
4. Wu, R.R., et al. 1997. cDNA cloning of the basement membrane chondroitin sulfate proteoglycan core protein, Bamacan: a five domain structure including coiled-coil motifs. *J. Cell Biol.* 136: 433-444.
5. Schmiesing, J.A., et al. 1998. Identification of two distinct human SMC protein complexes involved in mitotic chromosome dynamics. *Proc. Natl. Acad. Sci. USA* 95: 12906-12911.
6. Strunnikov, A.V., et al. 1999. Structural maintenance of chromosomes (SMC) proteins: conserved molecular properties for multiple biological functions. *Eur. J. Biochem.* 263: 6-13.
7. Nishiwaki, T., et al. 1999. Isolation and characterization of a human cDNA homologous to the *Xenopus laevis* XCAP-C gene belonging to the structural maintenance of chromosomes (SMC) family. *J. Hum. Genet.* 4: 197-202.
8. Ghiselli, G., et al. 1999. Complete cDNA cloning, genomic organization, chromosomal assignment, functional characterization of the promoter, and expression of the murine Bamacan gene. *J. Biol. Chem.* 274: 17384-17393.
9. Ghiselli, G., et al. 2000. Overexpression of bamacan/SMC3 causes transformation. *J. Biol. Chem.* 275: 20235-20238.

CHROMOSOMAL LOCATION

Genetic locus: SMC1B (human) mapping to 22q13.31; Smc1b (mouse) mapping to 15 E2.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

SOURCE

SMC1 β (C-16) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of SMC1 β of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-27961 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

SMC1 β (C-16) is recommended for detection of SMC1 β of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

SMC1 β (C-16) is also recommended for detection of SMC1 β in additional species, including bovine.

Suitable for use as control antibody for SMC1 β siRNA (h): sc-63348, SMC1 β siRNA (m): sc-38388, SMC1 β shRNA Plasmid (h): sc-63348-SH, SMC1 β shRNA Plasmid (m): sc-38388-SH, SMC1 β shRNA (h) Lentiviral Particles: sc-63348-V and SMC1 β shRNA (m) Lentiviral Particles: sc-38388-V.

Molecular Weight of SMC1 β : 150 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200 or NIH/3T3 whole cell lysate: sc-2210.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker[™] compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker[™] Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz[™] Mounting Medium: sc-24941.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

MONOS
Satisfaction
Guaranteed

Try **SMC1 (C2M): sc-56776**, our highly recommended monoclonal alternative to SMC1 β (C-16).