

# SDHA (C-18): sc-27992

## BACKGROUND

In aerobic respiration reactions, succinate dehydrogenase (SDH) catalyzes the oxidation of succinate and ubiquinone to fumarate and ubiquinol. Four subunits comprise the SDH protein complex: a flavochrome subunit (SDHA), an iron-sulfur protein (SDHB), and two membrane-bound subunits (SDHC and SDHD) anchored to the inner mitochondrial membrane. Mutations to these subunits cause mitochondrial dysfunction, corresponding to several distinct disorders. Mutations in the membrane bound components may cause hereditary paraganglioma, while SDHA mutations are associated with juvenile encephalopathy as well as Leigh Syndrome, a severe neurological disorder. Inactivating mutations in SDHB correlate with inherited, but not necessarily sporadic, cases of pheochromocytoma.

## CHROMOSOMAL LOCATION

Genetic locus: SDHA (human) mapping to 5p15.33; Sdha (mouse) mapping to 13 C1.

## SOURCE

SDHA (C-18) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the C-terminus of SDHA of human origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-27992 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## APPLICATIONS

SDHA (C-18) is recommended for detection of precursor and mature SDHA of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

SDHA (C-18) is also recommended for detection of precursor and mature SDHA in additional species, including canine, bovine and porcine.

Suitable for use as control antibody for SDHA siRNA (h): sc-61834, SDHA siRNA (m): sc-61835, SDHA shRNA Plasmid (h): sc-61834-SH, SDHA shRNA Plasmid (m): sc-61835-SH, SDHA shRNA (h) Lentiviral Particles: sc-61834-V and SDHA shRNA (m) Lentiviral Particles: sc-61835-V.


Molecular Weight of SDHA: 70 kDa.

Positive Controls: human heart tissue extract, mouse brain extract: sc-2253 or mouse heart extract: sc-2254.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## DATA



SDHA (C-18): sc-27992. Western blot analysis of SDHA expression in human heart (A), mouse heart (B) and mouse brain (C) tissue extracts.

## SELECT PRODUCT CITATIONS

- Fisher, K.W., et al. 2011. Kinase suppressor of ras 1 (KSR1) regulates PGC1 $\alpha$  and estrogen-related receptor  $\alpha$  to promote oncogenic Ras-dependent anchorage-independent growth. *Mol. Cell. Biol.* 31: 2453-2461.
- Deldicque, L., et al. 2012. Endurance training in mice increases the unfolded protein response induced by a high-fat diet. *J. Physiol. Biochem.* E-published.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.


 MONOS  
Satisfation  
Guaranteed

Try **SDHA (F-2): sc-390381** or **SDHA (D-4): sc-166947**, our highly recommended monoclonal alternatives to SDHA (C-18).