

MAT II α (I-17): sc-28031

BACKGROUND

Methionine adenosyltransferase (MAT) catalyzes the formation of S-adenosyltransferase (AdoMet) for methionine catabolism in the liver. Two different genes, MAT1A and MAT2A, encode a liver specific and non-liver specific form of MAT, respectively. Inactivation of the liver specific gene product, designated MAT I/III, associates with liver diseases such as cirrhosis. MAT1A expression also correlates with a differentiated phenotype, whereas liver cells expressing MAT2A present a dedifferentiated phenotype and lowered AdoMet synthesis. Likewise, NF κ B and TNF α cause a switch from MAT1A to MAT2A expression in human hepatocellular carcinoma (HCC), which facilitates cancer cell growth.

REFERENCES

1. Lu, S.C., et al. 2002. Role of abnormal methionine metabolism in alcoholic liver injury. *Alcohol* 27: 155-162.
2. Avila, M.A., et al. 2002. S-adenosylmethionine revisited: its essential role in the regulation of liver function. *Alcohol* 27: 163-167.
3. Martinez-Chantar, M.L., et al. 2003. L-methionine availability regulates expression of the methionine adenosyltransferase 2A gene in human hepatocarcinoma cells: role of S-adenosylmethionine. *J. Biol. Chem.* 278: 19885-19890.

CHROMOSOMAL LOCATION

Genetic locus: MAT2A (human) mapping to 2p11.2; Mat2a (mouse) mapping to 6 C1.

SOURCE

MAT II α (I-17) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of MAT II α of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-28031 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

MAT II α (I-17) is recommended for detection of MAT II α of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

MAT II α (I-17) is also recommended for detection of MAT II α in additional species, including equine, canine, bovine and porcine.


Molecular Weight of MAT II α : 44 kDa.

Positive Controls: Hep G2 cell lysate: sc-2227.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 4) Immunohistochemistry: use ImmunoCruz™: sc-2053 or ABC: sc-2023 goat IgG Staining Systems.

DATA



MAT II α (I-17): sc-28031. Western blot analysis of MAT II α expression in Hep G2 whole cell lysate.



MAT II α (I-17): sc-28031. Immunoperoxidase staining of formalin fixed, paraffin-embedded human kidney tissue showing cytoplasmic and nuclear staining of cells in glomeruli and cells in tubules.

SELECT PRODUCT CITATIONS

1. Brown, J.M., et al. 2010. Temporal study of acetaminophen (APAP) and S-adenosyl-L-methionine (SAME) effects on subcellular hepatic SAME levels and methionine adenosyltransferase (MAT) expression and activity. *Toxicol. Appl. Pharmacol.* 247: 1-9.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS
Satisfaction
Guaranteed

Try **MAT I α /II α (B-10): sc-166452** or **MAT II (F-12): sc-398917**, our highly recommended monoclonal alternatives to MAT II α (I-17).