

galectin-2 (H-45): sc-28249

BACKGROUND

Galectins are a family of soluble β -galactoside-binding animal lectins that modulate cell-to-cell adhesion and cell-to-extracellular matrix (ECM) interactions and play a role in tumor progression, pre-mRNA splicing and apoptosis. Galectin-2, also known as LGALS2, Lactose-binding lectin 2 or HL14, is structurally closely related to galectin-1, but is expressed primarily in the gastrointestinal tract. Galectin-2 induces apoptosis in activated T cells and binds to the cytokine lymphotoxin-alpha (LTA) with possible implications in risk of myocardial infarction.

REFERENCES

1. Couraud, P.O., et al. 1989. Molecular cloning, characterization, and expression of a human 14 kDa lectin. *J. Biol. Chem.* 264: 1310-1316.
2. Hirabayashi, J., et al. 1989. Cloning and nucleotide sequence of a full-length cDNA for human 14 kDa beta-galactoside-binding lectin. *Biochim. Biophys. Acta.* 1008: 85-91.
3. Mehrabian, M., et al. 1993. Two members of the S-lac lectin gene family, LGALS1 and LGALS2, reside in close proximity on human chromosome 22q12-q13. *Genomics.* 15: 418-420.
4. Sturm, A., et al. 2004. Human galectin-2: novel inducer of T cell apoptosis with distinct profile of caspase activation. *J. Immunol.* 173: 3825-3837.
5. Ozaki, K., et al. 2004. Functional variation in LGALS2 confers risk of myocardial infarction and regulates lymphotoxin-alpha secretion *in vitro*. *Nature* 429: 72-75.
6. Abedin, M.J., et al. 2003. Potential roles of galectins in myeloid differentiation into three different lineages. *J. Leukoc. Biol.* 73: 650-656.
7. Oka, T., et al. 1999. Identification and cloning of rat galectin-2: expression is predominantly in epithelial cells of the stomach. *Arch. Biochem. Biophys.* 361: 195-201.
8. Cornillot, J.D., et al. 1998. Production and characterization of a monoclonal antibody able to discriminate galectin-1 from galectin-2 and galectin-3. *Glycobiology* 8: 425-432.

CHROMOSOMAL LOCATION

Genetic locus: LGALS2 (human) mapping to 22q13.1.

SOURCE

galectin-2 (H-45) is a rabbit polyclonal antibody raised against amino acids 1-45 mapping at the N-terminus of galectin-2 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

galectin-2 (H-45) is recommended for detection of galectin-2 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


Suitable for use as control antibody for galectin-2 siRNA (h): sc-44532, galectin-2 shRNA Plasmid (h): sc-44532-SH and galectin-2 shRNA (h) Lentiviral Particles: sc-44532-V.

Positive Controls: galectin-2 (h2): 293T Lysate: sc-116773 or HISM cell lysate: sc-2229.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/ 2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



galectin-2 (H-45): sc-28249. Western blot analysis of galectin-2 expression in non-transfected: sc-117752 (A) and human galectin-2 transfected: sc-116773 (B) 293T whole cell lysates.



galectin-2 (H-45): sc-28249. Western blot analysis of galectin-2 expression in HISM whole cell lysate.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.