Ob (B-4): sc-28344



The Power to Question

BACKGROUND

Although there is substantial evidence that body weight is physiologically regulated, the molecular basis of obesity is unknown. Five single-gene mutations in mice that result in an obese phenotype have been identified. The first such recessive obesity mutation, the obese mutation (0b), was identified in 1950. Mutation of 0b (also designated leptin) results in profound obesity and type II diabetes as part of a syndrome that resembles morbid obesity in humans. It have been postulated that the 0b gene product may function as a component of a signaling pathway in adipose tissue that functions to regulate body fat depot size. The cloning and sequence analysis of the mouse 0b gene and its human homolog has been described. Ob encodes an adipose tissue-specific mRNA with a highly conserved 167 amino acid open reading frame. The predicted amino acid sequence is 84% identical between human and mouse and has the features of a secreted protein. A nonsense mutation in codon 105 has been found in the original congenic C57BL/6J 0b/0b mouse strain.

REFERENCES

- 1. Friedman, J.M., et al. 1991. Molecular mapping of the mouse Ob mutation. Genomics 11: 1054-1062.
- 2. Friedman, J.M. and Leibel, R.L. 1992. Tackling a weighty problem. Cell 69: 217-220.
- 3. Rink, T.J. 1994. In search of a satiety factor. Nature 372: 406-407.

CHROMOSOMAL LOCATION

Genetic locus: LEP (human) mapping to 7q32.1.

SOURCE

Ob (B-4) is a mouse monoclonal antibody raised against amino acids 22-167 of Ob of human origin.

PRODUCT

Each vial contains 200 μg lgG_{2a} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Ob (B-4) is available conjugated to agarose (sc-28344 AC), 500 $\mu g/0.25$ ml agarose in 1 ml, for IP.

APPLICATIONS

Ob (B-4) is recommended for detection of Ob of human origin by Western Blotting (starting dilution 1:100, dilution range), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


Suitable for use as control antibody for Ob siRNA (h): sc-37189, Ob shRNA Plasmid (h): sc-37189-SH and Ob shRNA (h) Lentiviral Particles: sc-37189-V.

Molecular Weight of Ob: 16 kDa.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-lgG κ BP-HRP: sc-516102 or m-lgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker^M Molecular Weight Standards: sc-2035, UltraCruz* Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-lgG κ BP-FITC: sc-516140 or m-lgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz* Mounting Medium: sc-24941 or UltraCruz* Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-lgG κ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

DATA





Ob (B-4): sc-28344. Western blot analysis of human recombinant Ob.

Ob (B-4): sc-28344. Immunoperoxidase staining of formalin fixed, paraffin-embedded human breast tissue showing extracellular staining of connective tissue.

SELECT PRODUCT CITATIONS

- Bressan, E., et al. 2019. Metal nanoparticles released from dental implant surfaces: potential contribution to chronic inflammation and peri-implant bone loss. Materials 12: 2036.
- 2. Schon, S.B., et al. 2022. Obesity-related alterations in protein expression in human follicular fluid from women undergoing in vitro fertilization. F S Sci. 3: 331-339.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.



See **Ob (F-3): sc-48408** for Ob antibody conjugates, including AC, HRP, FITC, PE, and Alexa Fluor® 488, 546, 594, 647, 680 and 790.