

MEL-1B-R (G-20): sc-28453

BACKGROUND

The melatonin receptors, MEL-1A-R and MEL-1B-R, are members of the superfamily of guanine nucleotide-binding regulatory protein (G protein)-coupled receptors. The melatonin receptors are activated by the hormone melatonin (Mel), which is secreted by the pineal gland at night as part of the circadian clock. MEL-1A-R is thought to be involved in pacing the biological clock. Both MEL-1A-R and MEL-1B-R are implicated in controlling cellular growth in response to melatonin. MEL-1B-R is an integral membrane protein expressed in retina and, to a lesser extent, in brain and hippocampus. Functional studies of NIH/3T3 cells stably expressing the MEL-1B-R melatonin receptor indicate that it is coupled to inhibition of adenylyl cyclase.

REFERENCES

1. Reppert, S.M., et al. 1995. Molecular characterization of a second melatonin receptor expressed in human retina and brain: the MEL1B melatonin receptor. *Proc. Natl. Acad. Sci. USA* 92: 8734-8738.
2. Reppert, S.M., et al. 1996. Cloning of a melatonin-related receptor from human pituitary. *FEBS Lett.* 386: 219-224.
3. Brzezinski, A. 1997. Melatonin in humans. *N. Engl. J. Med.* 336: 186-195.
4. Niles, L.P., et al. 1999. Melatonin receptor mRNA expression in human granulosa cells. *Mol. Cell. Endocrinol.* 156: 107-110.
5. Ebisawa, T., et al. 2000. Genetic polymorphisms of human melatonin 1B receptor gene in circadian rhythm sleep disorders and controls. *Neurosci. Lett.* 280: 29-32.
6. Yuan, L., et al. 2002. MT₁ melatonin receptor overexpression enhances the growth suppressive effect of melatonin in human breast cancer cells. *Mol. Cell. Endocrinol.* 192: 147-156.

CHROMOSOMAL LOCATION

Genetic locus: MTNR1B (human) mapping to 11q14.3.

SOURCE

MEL-1B-R (G-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an extracellular domain of MEL-1B-R of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-28453 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

APPLICATIONS

MEL-1B-R (G-20) is recommended for detection of MEL-1B-R of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

MEL-1B-R (G-20) is also recommended for detection of MEL-1B-R in additional species, including porcine.

Suitable for use as control antibody for MEL-1B-R siRNA (h): sc-40114, MEL-1B-R shRNA Plasmid (h): sc-40114-SH and MEL-1B-R shRNA (h) Lentiviral Particles: sc-40114-V.


Molecular Weight of MEL-1B-R: 36 kDa.

Positive Controls: Y79 cell lysate: sc-2240, Jurkat whole cell lysate: sc-2204 or IMR-32 cell lysate: sc-2409.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



MEL-1B-R (G-20): sc-28453. Western blot analysis of MEL-1B-R expression in Y79 (A), Jurkat (B) and IMR-32 (C) whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.



Try **MEL-1A/B-R (B-8): sc-398788**, our highly recommended monoclonal alternative to MEL-1B-R (G-20).