

# GSTM (M-17)-R: sc-28504-R

## BACKGROUND

Members of the glutathione S-transferase (GST) family of proteins function in the detoxification of xenobiotics to protect cells against toxicant-induced damage. There are eight families of GST proteins, namely  $\alpha$ ,  $\kappa$ ,  $\mu$ ,  $\omega$ ,  $\pi$ ,  $\sigma$ ,  $\theta$  and  $\zeta$ , each of which are composed of proteins that have a variety of functions throughout the cell. The GSTM proteins (GSTM1-GSTM5 in human and GSTM1-GSTM7 in mouse) are members of the  $\mu$  class of enzymes that conjugate with glutathione and function in the detoxification of carcinogens, environmental toxins and products of oxidative stress. The  $\mu$  proteins are encoded by genes that form a cluster on chromosome 1p13.3 and are known to be highly polymorphic, affecting an individual's susceptibility to toxins. Mutations in the genes encoding GSTM proteins are associated with tumor progression, suggesting an important role for GSTM proteins in carcinogenesis.

## REFERENCES

- McGuire, S., et al. 1997. Increased levels of glutathione S transferases and appearance of novel  $\alpha$  class isoenzymes in kidneys of mice exposed to mercuric chloride. I. Biochemical and immunohistochemical studies. *Nephron* 77: 452-460.
- Massey, T.E., et al. 2000. Mechanisms of Aflatoxin B1 lung tumorigenesis. *Exp. Lung Res.* 26: 673-683.
- Raza, H., et al. 2002. Multiple isoforms of mitochondrial glutathione S-transferases and their differential induction under oxidative stress. *Biochem. J.* 366: 45-55.
- Bartley, P.A., et al. 2003. Regulation of the gene encoding glutathione S-transferase M1 (GSTM1) by the Myb oncoprotein. *Oncogene* 22: 7570-7575.
- Breton, C.V., et al. 2007. GSTM1 and APE1 genotypes affect arsenic-induced oxidative stress: a repeated measures study. *Environ Health.* 6: 39.
- Shang, W., et al. 2008. Expressions of glutathione S-transferase  $\alpha$ ,  $\mu$ , and  $\pi$  in brains of medically intractable epileptic patients. *BMC Neurosci.* 9: 67.
- Lucena, M.I., et al. 2008. Glutathione S-transferase M1 and T1 null genotypes increase susceptibility to idiosyncratic drug-induced liver injury. *Hepatology* 48: 588-596.
- Fukino, K., et al. 2008. Effects of N-acetyltransferase 2 (NAT2), CYP2E1 and glutathione-S-transferase (GST) genotypes on the serum concentrations of isoniazid and metabolites in tuberculosis patients. *J. Toxicol. Sci.* 33: 187-195.
- Agúndez, J.A., et al. 2008. Glutathione S-transferase GSTT1 and GSTM1 allozymes: beyond null alleles. *Pharmacogenomics* 9: 359-363.

## SOURCE

GSTM (M-17)-R is an affinity purified rabbit polyclonal antibody raised against a peptide mapping near the C-terminus of GSTM1 of mouse origin.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PRODUCT

Each vial contains 200  $\mu$ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-28504 P, (100  $\mu$ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## APPLICATIONS

GSTM (M-17)-R is recommended for detection of GSTM1, GSTM6 and GSTM7 of mouse and rat origin and, to a lesser extent, GSTM1, GSTM2 and GSTM4 of human origin and GSTM2 of mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

GSTM (M-17)-R is also recommended for detection of GSTM proteins in additional species, including equine.

Molecular Weight of all GSTM proteins: 26 kDa.

Positive Controls: 3T3-L1 cell lysate: sc-2243 or mouse liver extract: sc-2256.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.



Try **GSTM1 (1H4F2): sc-517197**, our highly recommended monoclonal alternative to GSTM (M-17).