

Cdc28 (γ-40): sc-28550

BACKGROUND

Cell cycle progression is controlled at a point late in G₁ designated Start. Passage through Start requires the activity of the cyclin-dependent protein kinase Cdc28. Transition from G₁ to S phase requires the association of Cdc28 with members of the G₁ cyclin family, including Cln1, Cln2 and Cln3 (also designated DAF-1 or WHI1). The G₂ to M phase requires the M phase cyclins, Clb1 (also designated Scb1) and Clb2, as well as the G₂ cyclins, Clb3 and Clb4. The S phase cyclins Clb5 and Clb6 coordinate DNA replication with cytokinesis. Expression of the cyclins is controlled by Ubc9 and Cdc34 (also designated Udc3 or Dna6) via ubiquitin-mediated proteolysis.

REFERENCES

- Nasmyth, K. 1993. Control of the yeast cell cycle by the Cdc28 protein kinase. *Curr. Opin. Cell Biol.* 5: 166-179.
- Sherlock, G. and Rosamond, J. 1993. Starting to cycle: G₁ controls regulating cell division in budding yeast. *J. Gen. Microbiol.* 139: 2531-2541.
- Amon, A., Tyers, M., Futcher, B. and Nasmyth, K. 1993. Mechanisms that help the yeast cell cycle clock tick: G₂ cyclins transcriptionally activate G₂ cyclins and repress G₁ cyclins. *Cell* 74: 993-1007.
- Basco, R.D., Segal, M.D. and Reed, S.I. 1995. Negative regulation of G₁ and G₂ by S-phase cyclins of *Saccharomyces cerevisiae*. *Mol. Cell. Biol.* 15: 5030-5042.
- Seufert, W., Futcher, B. and Jentsch, S. 1995. Role of a ubiquitin-conjugating enzyme in degradation of S- and M- phase cyclins. *Nature* 373: 78-81.
- Prendergast, J.A., Ptak, C., Arnason, T.G. and Ellison, M.J. 1995. Increased ubiquitin expression suppresses the cell cycle defect associated with the yeast ubiquitin conjugating enzyme, Cdc34 (Ubc3). Evidence for a noncovalent interaction between Cdc34 and ubiquitin. *J. Biol. Chem.* 270: 9347-9352.
- Levine, K., Huang, K. and Cross, F.R. 1996. *Saccharomyces cerevisiae* G₁ cyclins differ in their intrinsic functional specificities. *Mol. Cell. Biol.* 16: 6794-6803.
- Blondel, M. and Mann, C. 1996. G₂ cyclins are required for the degradation of G₁ cyclins in yeast. *Nature* 384: 279-282.

SOURCE

Cdc28 (γ-40) is a rabbit polyclonal antibody raised against amino acids 91-130 mapping within an internal region of Cdc28 of *Saccharomyces cerevisiae* origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.


APPLICATIONS

Cdc28 (γ-40) is recommended for detection of Cdc28 of *Saccharomyces cerevisiae* origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



Cdc28 (γ-40): sc-28550. Western blot analysis of yeast recombinant Cdc28 fusion protein.

SELECT PRODUCT CITATIONS

- Keating, P., Rachidi, N., Tanaka, T.U. and Stark, M.J. 2009. Ipl1-dependent phosphorylation of Dam1 is reduced by tension applied on kinetochores. *J. Cell Sci.* 122: 4375-4382.
- Rachfall, N., Heinemeyer, I., Morgenstern, B., Valerius, O. and Braus, G.H. 2011. 5'TRU: identification and analysis of translationally regulative 5'untranslated regions in amino acid starved yeast cells. *Mol. Cell Proteomics* 10: M110.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

MONOS
Satisfaction
Guaranteed

Try **Cdc28 (G-7): sc-515762**, our highly recommended monoclonal alternative to Cdc28 (γ-40).