

## KCNE3 (H-80): sc-28796

### BACKGROUND

Voltage-gated K<sup>+</sup> channels in the plasma membrane control the repolarization and the frequency of action potentials in neurons, muscles, and other excitable cells. KCNE3 (potassium voltage-gated channel, Isk-related family, member 3), also known as HYPK, potassium channel subunit  $\beta$  MiRP2 or HOKPP, is a 103 amino acid single-pass type I membrane protein belonging to the potassium channel KCNE family. Expressed predominantly in kidney, KCNE3 is also found at moderate levels in small intestine and associates with a voltage-gated potassium channel complex to regulate stability and gating kinetics. The gene encoding KCNE3 maps to human chromosome 11q13.4; defects in which are the cause of an autosomal dominant disorder known as periodic paralysis hypokalemic (or HOKPP), a muscular disorder known as thyrotoxic periodic paralysis type 1 (TTPP1) and Brugada syndrome type 6 (BRS6).

### REFERENCES

- Abbott, G.W., et al. 1999. MiRP1 forms IKr potassium channels with HERG and is associated with cardiac arrhythmia. *Cell* 97: 175-187.
- Abbott, G.W., et al. 2001. MiRP2 forms potassium channels in skeletal muscle with Kv3.4 and is associated with periodic paralysis. *Cell* 104: 217-231.
- Dias Da Silva, M.R., et al. 2002. A mutation in the KCNE3 potassium channel gene is associated with susceptibility to thyrotoxic hypokalemic periodic paralysis. *J. Clin. Endocrinol. Metab.* 87: 4881-4884.
- Tang, N.L., et al. 2004. No mutation in the KCNE3 potassium channel gene in Chinese thyrotoxic hypokalaemic periodic paralysis patients. *Clin. Endocrinol.* 61: 109-112.
- Lundby, A. and Olesen, S.P. 2006. KCNE3 is an inhibitory subunit of the Kv4.3 potassium channel. *Biochem. Biophys. Res. Commun.* 346: 958-967.
- Abbott, G.W., et al. 2006. Phosphorylation and protonation of neighboring MiRP2 sites: function and pathophysiology of MiRP2-Kv3.4 potassium channels in periodic paralysis. *FASEB J.* 20: 293-301.

### CHROMOSOMAL LOCATION

Genetic locus: KCNE3 (human) mapping to 11q13.4; Kcne3 (mouse) mapping to 7 E2.

### SOURCE

KCNE3 (H-80) is a rabbit polyclonal antibody raised against amino acids 1-80 mapping at the N-terminus of KCNE3 of human origin.

### PRODUCT

Each vial contains 200  $\mu$ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

### RESEARCH USE

For research use only, not for use in diagnostic procedures.

### APPLICATIONS

KCNE3 (H-80) is recommended for detection of KCNE3 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

KCNE3 (H-80) is also recommended for detection of KCNE3 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for KCNE3 siRNA (h): sc-35743, KCNE3 siRNA (m): sc-35744, KCNE3 shRNA Plasmid (h): sc-35743-SH, KCNE3 shRNA Plasmid (m): sc-35744-SH, KCNE3 shRNA (h) Lentiviral Particles: sc-35743-V and KCNE3 shRNA (m) Lentiviral Particles: sc-35744-V.


Molecular Weight of KCNE3: 12 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200.

### RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

### DATA



KCNE3 (H-80): sc-28796. Western blot analysis of human recombinant KCNE3 fusion protein.

### STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

**MONOS**  
Satisfaction  
Guaranteed

Try **KCNE3 (G-6): sc-393841**, our highly recommended monoclonal alternative to KCNE3 (H-80).