

# GluR-5 (H-50): sc-28798

## BACKGROUND

Glutamate receptors mediate most excitatory neurotransmission in the brain and play an important role in neural plasticity, neural development and neurodegeneration. Ionotropic glutamate receptors are categorized into NMDA receptors and kainate/AMPA receptors, both of which contain glutamate-gated, cation-specific ion channels. Kainate/AMPA receptors are co-localized with NMDA receptors in many synapses and consist of seven structurally related subunits designated GluR-1 to -7. The kainate/AMPA receptors are primarily responsible for the fast excitatory neurotransmission by glutamate, whereas the NMDA receptors are functionally characterized by a slow kinetic and a high permeability for Ca<sup>2+</sup> ions. The NMDA receptors consist of five subunits:  $\epsilon$  1, 2, 3, 4 and one  $\zeta$  subunit. The  $\zeta$  subunit is expressed throughout the brainstem, whereas the four  $\epsilon$  subunits display limited distribution.

## REFERENCES

- Choi, D.W., et al. 1990. The role of glutamate neurotoxicity in hypoxic-ischemic neuronal death. *Annu. Rev. Neurosci.* 13: 171-182.
- Nakanishi, S. 1992. Molecular diversity of glutamate receptors and implications for brain function. *Science* 258: 597-603.
- Stern, P., et al. 1992. Fast and slow components of unitary EPSCs on stellate cells elicited by focal stimulation in slices of rat visual cortex. *J. Physiol.* 449: 247-278.
- Bliss, T.V., et al. 1993. A synaptic model of memory: long-term potentiation in the hippocampus. *Nature* 361: 31-39.

## CHROMOSOMAL LOCATION

Genetic locus: GRIK1 (human) mapping to 21q21.3.

## SOURCE

GluR-5 (H-50) is a rabbit polyclonal antibody raised against amino acids 869-918 mapping at the C-terminus of GluR-5 of human origin.

## PRODUCT

Each vial contains 200  $\mu$ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## APPLICATIONS

GluR-5 (H-50) is recommended for detection of GluR-5 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

GluR-5 (H-50) is also recommended for detection of GluR-5 in additional species, including equine, canine, porcine and avian.


Suitable for use as control antibody for GluR-5 siRNA (h): sc-42487, GluR-5 shRNA Plasmid (h): sc-42487-SH and GluR-5 shRNA (h) Lentiviral Particles: sc-42487-V.

Molecular Weight of GluR-5: 105-110 kDa.


## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## DATA



GluR-5 (H-50): sc-28798. Western blot analysis of human recombinant GluR-5 fusion protein.



GluR-5 (H-50): sc-28798. Western blot analysis of GluR-5 expression in 293T whole cell lysate.

## SELECT PRODUCT CITATIONS

- Zhang, H.J., et al. 2012. ATPA induced GluR5-containing kainite receptor S-nitrosylation via activation of GluR5-Gq-PLC-IP(3)R pathway and signalling module GluR5-PSD-95-nNOS. *Int. J. Biochem. Cell Biol.* 44: 2261-2271.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.



Try **GluR-5 (E-12): sc-393420**, our highly recommended monoclonal alternative to GluR-5 (H-50).