

eIF2B γ (H-300): sc-28853

BACKGROUND

The initiation of protein synthesis in eukaryotic cells is regulated by interactions between protein initiation factors and RNA molecules. The eukaryotic initiation complex eIF2B exists as a five subunit complex composed of eIF2B α , eIF2B β , eIF2B γ , eIF2B δ and eIF2B ϵ . The eIF2B complex catalyzes the exchange of GDP for GTP on the eIF2 complex, following the interaction of eIF2/GTP with the 40S ribosomal subunit. Guanine nucleotide exchange factor (GEF) activity was exhibited by the eIF2B ϵ subunit alone, but it was greater in the presence of all five eIF2B subunits. Phosphorylation of eIF2 inhibits GEF activity of eIF2B, an inhibition that requires the eIF2B α subunit.

REFERENCES

- Henderson, R.A., et al. 1994. The δ -subunit of murine guanine nucleotide exchange factor eIF-2B. Characterization of cDNAs predicts isoforms differing at the amino-terminal end. *J. Biol. Chem.* 269: 30517-30523.
- Flowers, K.M., et al. 1995. Structure and sequence of the gene encoding the α -subunit of rat translation initiation factor-2B. *Biochim. Biophys. Acta* 1264: 163-167.
- Price, N.T., et al. 1996. Cloning of cDNA for the γ -subunit of mammalian translation initiation factor 2B, the guanine nucleotide-exchange factor for eukaryotic initiation factor 2. *Biochem. J.* 318: 631-636.
- Price, N.T., et al. 1996. eIF2B, the guanine nucleotide-exchange factor for eukaryotic initiation factor 2. Sequence conservation between the α , β and δ subunits of eIF2B from mammals and yeast. *Biochem. J.* 318: 637-643.
- Asuru, A.I., et al. 1996. Cloning and characterization of cDNAs encoding the ϵ -subunit of eukaryotic initiation factor-2B from rabbit and human. *Biochim. Biophys. Acta* 1307: 309-317.
- Webb, B.L. and Proud, C.G. 1997. Eukaryotic initiation factor 2B (eIF2B). *Int. J. Biochem. Cell Biol.* 29: 1127-1131.

CHROMOSOMAL LOCATION

Genetic locus: EIF2B3 (human) mapping to 1p34.1; Eif2b3 (mouse) mapping to 4 D1.

SOURCE

eIF2B γ (H-300) is a rabbit polyclonal antibody raised against amino acids 153-452 mapping at the C-terminus of eIF2B γ of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4 $^{\circ}$ C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

APPLICATIONS

eIF2B γ (H-300) is recommended for detection of eIF2B γ of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


eIF2B γ (H-300) is also recommended for detection of eIF2B γ in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for eIF2B γ siRNA (h): sc-35274, eIF2B γ siRNA (m): sc-35275, eIF2B γ shRNA Plasmid (h): sc-35274-SH, eIF2B γ shRNA Plasmid (m): sc-35275-SH, eIF2B γ shRNA (h) Lentiviral Particles: sc-35274-V and eIF2B γ shRNA (m) Lentiviral Particles: sc-35275-V.

Molecular Weight of eIF2B γ : 50 kDa.

Positive Controls: NIH/3T3 nuclear extract: sc-2138, K-562 nuclear extract: sc-2130 or HeLa nuclear extract: sc-2120.

DATA



eIF2B γ (H-300): sc-28853. Western blot analysis of eIF2B γ expression in HeLa nuclear extract.

SELECT PRODUCT CITATIONS

- Martin, L., et al. 2010. Regulation of the unfolded protein response by eIF2B δ isoforms. *J. Biol. Chem.* 285: 31944-31953.

RESEARCH USE

For research use only, not for use in diagnostic procedures.



Try **eIF2B γ (F-7): sc-514230** or **eIF2B γ (P-5): sc-9980**, our highly recommended monoclonal alternatives to eIF2B γ (H-300).