

TFPI (H-120): sc-28861

BACKGROUND

The extrinsic pathway of blood coagulation is initiated by contact of plasma factor VII with tissue factor, a cellular membrane glycoprotein that normally is segregated from the bloodstream but can be exposed after tissue injury or newly synthesized in endothelial cells or leukocytes after stimulation by endotoxin and cytokines. Inhibition of factor VIIa/tissue factor activity requires a plasma component (tissue factor pathway inhibitor (TFPI), lipoprotein-associated coagulation inhibitor (LACI) or extrinsic pathway inhibitor (EPI)) and factor Xa. TFPI directly inhibits factor Xa, and, in an Xa-dependent fashion, also inhibits the factor VIIa-tissue factor catalytic complex. TFPI is a multivalent, Kunitz-type proteinase inhibitor that circulates in association with plasma lipoproteins VLDL, LDL, and HDL. TFPI-2 (also known as placental protein 5) is a related glycoprotein that was originally isolated from human placenta.

REFERENCES

1. Broze, G.J., Jr. and Miletich, J.P. 1987. Characterization of the inhibition of tissue factor in serum. *Blood* 69: 150-155.
2. Rao, L.V., et al. 1987. Studies of a mechanism inhibiting the initiation of the extrinsic pathway of coagulation. *Blood* 69: 645-651.
3. Davie, E.W., et al. 1991. The coagulation cascade: initiation, maintenance, and regulation. *Biochemistry* 30: 10363-10370.
4. Girard, T.J., et al. 1991. Structure of the human lipoprotein-associated coagulation inhibitor gene. Intro/exon gene organization and localization of the gene to chromosome 2. *J. Biol. Chem.* 266: 5036-5041.
5. Enyoji, K., et al. 1993. Human tissue factor pathway inhibitor (TFPI) gene: complete genomic structure and localization on the genetic map of chromosome 2q. *Genomics* 17: 423-428.
6. Kisiel, W., et al. 1994. Evidence that a second human tissue factor pathway inhibitor (TFPI-2) and human placental protein 5 are equivalent. *Blood* 84: 4384-4385.
7. Miyagi, Y., et al. 1996. Assignment of the human PP5/TFPI-2 gene to 7q22 by FISH and PCR-based human/rodent cell hybrid mapping panel analysis. *Genomics* 35: 267-268.

CHROMOSOMAL LOCATION

Genetic locus: TFPI (human) mapping to 2q32.1; Tfpi (mouse) mapping to 2 D.

SOURCE

TFPI (H-120) is a rabbit polyclonal antibody raised against amino acids 31-150 mapping near the N-terminus of TFPI of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

TFPI (H-120) is recommended for detection of precursor and mature forms of TFPI of human and, to a lesser extent, mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for TFPI siRNA (h): sc-41060, TFPI siRNA (m): sc-41061, TFPI shRNA Plasmid (h): sc-41060-SH, TFPI shRNA Plasmid (m): sc-41061-SH, TFPI shRNA (h) Lentiviral Particles: sc-41060-V and TFPI shRNA (m) Lentiviral Particles: sc-41061-V.


Molecular Weight of TFPI: 40 kDa.

Positive Controls: JAR cell lysate: sc-2276, K-562 whole cell lysate: sc-2203 or HeLa whole cell lysate: sc-2200.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



TFPI (H-120): sc-28861. Western blot analysis of TFPI expression in K-562 whole cell lysate.

RESEARCH USE

For research use only, not for use in diagnostic procedures.



Try **TFPI (G-5): sc-365920** or **TFPI (G-6): sc-133139**, our highly recommended monoclonal alternatives to TFPI (H-120).