

HIPK1 (H-50): sc-28896

BACKGROUND

The Homeodomain-Interacting Protein Kinase (HIPK) family, which includes HIPK1, HIPK2, HIPK3, contains a conserved protein kinase domain as well as a separate domain that interacts with homeoproteins. HIPK2, the most highly characterized family member, is thought to act as a co-repressor of homeodomain transcription factors as HIPK2 has been shown to enhance the DNA binding of the NK-3 homeoprotein *in vitro*. It is regulated by a posttranslational modification of a ubiquitin-like protein, SUMO-1, via covalent bonding to a lysine residue on HIPK2. This is similar to the binding of SUMO-1 to PML and Sp100. The conjugation of SUMO-1 is thought to direct each of these proteins to nuclear bodies (NB's), which appear to play a role in auto-immunity and viral protection. HIPK2 is the first protein kinase to be directed to nuclear bodies in response to ubiquitin-like modification.

REFERENCES

- Kim, Y.H., et al. 1998. Homeodomain-interacting Protein Kinases, a novel family of co-repressors for homeodomain transcription factors. *J. Biol. Chem.* 273: 25875-25879.
- Schwarz, S.E., et al. 1998. The ubiquitin-like proteins SMT3 and SUMO-1 are conjugated by the UBC9 E2 enzyme. *Proc. Natl. Acad. Sci. USA* 95: 560-564.
- Kim, Y.H., et al. 1999. Covalent modification of the homeodomain-interacting protein kinase 2 (HIPK2) by the ubiquitin-like protein SUMO-1. *Proc. Natl. Acad. Sci. USA* 96: 12350-12355.
- Muller, S. and Dejean, A. 1999. Viral immediate-early proteins abrogate the modification of SUMO-1 of PML and Sp100 proteins, correlating with nuclear body disruption. *J. Virol.* 73: 5137-5143.
- Ishov, A.M., et al. 1999. PML is critical for ND10 formation and recruits the PML-interacting protein daxx to this nuclear structure when modified by SUMO-1. *J. Cell. Biol.* 147: 221-234.
- Sternsdorf, T., et al. 1999. The nuclear dot protein sp100, characterization of domains necessary for dimerization, subcellular localization, and modification by small ubiquitin-like modifiers. *J. Biol. Chem.* 274: 12555-12566.

CHROMOSOMAL LOCATION

Genetic locus: HIPK1 (human) mapping to 1p13.2; Hipk1 (mouse) mapping to 3 F2.2.

SOURCE

HIPK1 (H-50) is a rabbit polyclonal antibody raised against amino acids 31-80 mapping near the N-terminus of HIPK1 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

HIPK1 (H-50) is recommended for detection of HIPK1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

HIPK1 (H-50) is also recommended for detection of HIPK1 in additional species, including equine, canine and bovine.

Suitable for use as control antibody for HIPK1 siRNA (h): sc-39048, HIPK1 siRNA (m): sc-39049, HIPK1 shRNA Plasmid (h): sc-39048-SH, HIPK1 shRNA Plasmid (m): sc-39049-SH, HIPK1 shRNA (h) Lentiviral Particles: sc-39048-V and HIPK1 shRNA (m) Lentiviral Particles: sc-39049-V.

Molecular Weight of HIPK1: 120 kDa.

Positive Controls: DU 145 cell lysate: sc-2268, MCF7 whole cell lysate: sc-2206 or SW480 cell lysate: sc-2219.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.


 MONOS
Satisfaction
Guaranteed

Try **HIPK1 (SR-5): sc-100382**, our highly recommended monoclonal alternative to HIPK1 (H-50).