

# GRIP1 (R-91): sc-28934

## BACKGROUND

Glutamate receptors mediate most excitatory neurotransmission in the brain and play an important role in neural plasticity, neural development and neurodegeneration. The glutamate receptor interacting proteins, GRIP1 and GRIP2, are members of the PDZ domain-containing protein family, and they specifically bind to the carboxy-terminus of AMPA receptor subunits, GluR-2 and GluR-3. GRIP1 and GRIP2 are involved in the targeting of GluR-2 and GluR-3 to the synapse. GRIP1 and GRIP2 are widely expressed in brain, with the highest levels in the cerebral cortex, hippocampus and olfactory bulb. They are both enriched in synaptic plasma and postsynaptic density fractions. GRIP1 is expressed in early development before the expression of AMPA receptors, specifically postnatal days 8-10, while GRIP2 expression parallels that of AMPA receptors during later developmental stages. GRIP1 and GRIP2 may mediate the endocytotic rate of GluR-2 and GluR-3 in response to the phosphorylation of the receptors on Ser 880 by PKC, which is implicated in the induction of cerebellar long-term depression (LTD).

## REFERENCES

1. Choi, D.W. and Rothman, S.M. 1990. The role of glutamate neurotoxicity in hypoxic-ischemic neuronal death. *Annu. Rev. Neurosci.* 13: 171-182.
2. Nakanishi, S. 1992. Molecular diversity of glutamate receptors and implications for brain function. *Science* 258: 597-603.
3. Wyszynski, M., Valtschanoff, J.G., Naisbitt, S., Dunah, A.W., Kim, E., Standaert, D.G., Weinberg, R. and Sheng, M. 1999. Association of AMPA receptors with a subset of glutamate receptor-interacting protein *in vivo*. *J. Neurosci.* 19: 6528-6537.
4. Dong, H., Zhang, P., Song, I., Petralia, R.S., Liao, D. and Huganir, R.L. 1999. Characterization of the glutamate receptor-interacting proteins GRIP1 and GRIP2. *J. Neurosci.* 19: 6930-6941.
5. Osten, P., Khatri, L., Perez, J.L., Kohr, G., Giese, G., Daly, C., Schulz, T.W., Wensky, A., Lee, L.M. and Ziff, E.B. 2000. Mutagenesis reveals a role for ABP/GRIP binding to GluR2 in synaptic surface accumulation of the AMPA receptor. *Neuron* 27: 313-325.
6. Matsuda, S., Launey, T., Mikawa, S. and Hirai, H. 2000. Disruption of AMPA receptor GluR-2 clusters following long-term depression induction in cerebellar Purkinje neurons. *EMBO J.* 19: 2765-2774.
7. Yamazaki, M., Fukaya, M., Abe, M., Ikeno, K., Kakizaki, T., Watanabe, M. and Sakimura, K. 2001. Differential palmitoylation of two mouse glutamate receptor interacting protein 1 forms with different N-terminal sequences. *Neurosci. Lett.* 304: 81-84.

## CHROMOSOMAL LOCATION

Genetic locus: GRIP1 (human) mapping to 12q14.3; Grip1 (mouse) mapping to 10 D2.

## SOURCE

GRIP1 (R-91) is a rabbit polyclonal antibody raised against amino acids 910-1100 mapping near the C-terminus of GRIP1 of rat origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## APPLICATIONS

GRIP1 (R-91) is recommended for detection of GRIP1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

GRIP1 (R-91) is also recommended for detection of GRIP1 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for GRIP1 siRNA (h): sc-42160, GRIP1 siRNA (m): sc-42161, GRIP1 shRNA Plasmid (h): sc-42160-SH, GRIP1 shRNA Plasmid (m): sc-42161-SH, GRIP1 shRNA (h) Lentiviral Particles: sc-42160-V and GRIP1 shRNA (m) Lentiviral Particles: sc-42161-V.

Molecular Weight of GRIP1: 130 kDa.

Positive Controls: IMR-32 cell lysate: sc-2409.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.

**MONOS**  
Satisfaction  
Guaranteed

Try **GRIP1 (H-4): sc-365937** or **GRIP1 (32): sc-135931**, our highly recommended monoclonal alternatives to GRIP1 (R-91).