

NK-2R (M-48): sc-28951

BACKGROUND

Substance P (SP) and neurokinin-A (NK-A) are members of the tachykinins, and they function as modulators of the immune and hematopoietic systems. The tachykinins interact with each of three cloned neurokinin (NK) receptors (NK-1R, NK-2R, NK-3R), with SP and NK-A exhibiting binding preferences for NK-1R and NK-2R, respectively. NK-4R shares close homology with NK-3R, and both have nearly identical pharmacological properties. In the normal ileum and colon, NK-1R and NK-2R are localized to smooth muscle cells of the muscularis mucosae and propria and to a few inflammatory cells of the lamina propria. NK-1R expression is also found in the muscular wall of submucosal blood vessels, enteric neurons and, to a lesser degree, in surface epithelial cells. NK-3R is found in the spinal cord in both lamina X and lamina II.

REFERENCES

1. Rameshwar, P. and Gascon, P. 1995. Substance P (SP) mediates production of stem cell factor and interleukin-1 in bone marrow stroma: potential autoregulatory role for these cytokines in SP receptor expression and induction. *Blood* 86: 482-490.
2. Rameshwar, P. and Gascon, P. 1997. Hematopoietic modulation by the tachykinins. *Acta Haematol.* 98: 59-64.
3. Zerari, F., Karpitskiy, V., Krause, J., Descarries, L. and Couture, R. 1997. Immunoelectron microscopic localization of NK-3 receptor in the rat spinal cord. *Neuroreport* 8: 2661-2664.
4. Sarau, H.M., Mooney, J.L., Schmidt, D.B., Foley, J.J., Buckley, P.T., Giardina, G.A., Wang, D.Y., Lee, J.A. and Hay, D.W. 2000. Evidence that the proposed novel human "neurokinin-4" receptor is pharmacologically similar to the human neurokinin-3 receptor but is not of human origin. *Mol. Pharmacol.* 58: 552-559.

CHROMOSOMAL LOCATION

Genetic locus: TACR2 (human) mapping to 10q22.1; Tacr2 (mouse) mapping to 10 B4.

SOURCE

NK-2R (M-48) is a rabbit polyclonal antibody raised against amino acids 207-254 mapping within an internal region of NK-2R of mouse origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

APPLICATIONS

NK-2R (M-48) is recommended for detection of NK-2R of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

NK-2R (M-48) is also recommended for detection of NK-2R in additional species, including bovine and porcine.


Suitable for use as control antibody for NK-2R siRNA (h): sc-42292, NK-2R siRNA (m): sc-42293, NK-2R shRNA Plasmid (h): sc-42292-SH, NK-2R shRNA Plasmid (m): sc-42293-SH, NK-2R shRNA (h) Lentiviral Particles: sc-42292-V and NK-2R shRNA (m) Lentiviral Particles: sc-42293-V.

Positive Controls: Jurkat whole cell lysate: sc-2204 or mouse liver extract: sc-2256.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 4) Immunohistochemistry: use ImmunoCruz™: sc-2051 or ABC: sc-2018 rabbit IgG Staining Systems.

DATA



NK-2R (M-48): sc-28951. Western blot analysis of NK-2R expression in Jurkat whole cell lysate (A) and mouse liver tissue extract (B).



NK-2R (M-48): sc-28951. Immunoperoxidase staining of formalin fixed, paraffin-embedded human heart muscle tissue showing cytoplasmic and membrane staining of myocytes.

RESEARCH USE

For research use only, not for use in diagnostic procedures.