

MC5-R (H-55): sc-28993

BACKGROUND

The melanocortin family comprises the α -, β - and γ - melanocyte stimulating hormones (MSH) and adrenocorticotrophin. The receptors for these hormones are seven-transmembrane, G protein-coupled proteins that activate adenyl cyclase. Five melanocortin receptors have been cloned and shown to exhibit different affinities and patterns of expression. MC1-R (MSH-R) is expressed in melanocytes and corticoadrenal tissue. MC2-R is the ACTH receptor and is expressed primarily in the adrenal cortex. MC3-R has been found in specific regions of the brain and is also expressed in placenta and gut. MC4-R is expressed primarily in brain, while MC5-R is expressed at low levels in most tissues.

REFERENCES

1. Mountjoy, K.G., et al. 1992. The cloning of a family of genes that encode the melanocortin receptors. *Science* 257: 1248-1251.
2. Chhajlani, V., et al. 1992. Molecular cloning and expression of the human melanocyte stimulating hormone receptor cDNA. *FEBS Lett.* 309: 417-420.
3. Gantz, I., et al. 1993. Molecular cloning of a novel melanocortin receptor. *J. Biol. Chem.* 268: 8246-8250.
4. Gantz, I., et al. 1993. Molecular cloning, expression and gene localization of a fourth melanocortin receptor. *J. Biol. Chem.* 268: 15174-15179.
5. Roselli-Rehffuss, L., et al. 1993. Identification of a receptor for γ -melanotropin and other pro-opiomelanocortin peptides in the hypothalamus and limbic system. *Proc. Natl. Acad. Sci. USA* 90: 8856-8860.
6. Chhajlani, V., et al. 1993. Molecular cloning of a novel human melanocortin receptor. *Biochem. Biophys. Res. Commun.* 195: 866-873.
7. Mountjoy, K.G., et al. 1994. Localization of the melanocortin 4 receptor (MC4-R) in neuroendocrine and autonomic control circuits in the brain. *Mol. Endocrinol.* 8: 1298-1308.
8. Labbe, O., et al. 1994. Molecular cloning of a mouse melanocortin 5 receptor gene widely expressed in peripheral tissues. *Biochemistry* 33: 4543-4549.

CHROMOSOMAL LOCATION

Genetic locus: MC5R (human) mapping to 18p11.21.

SOURCE

MC5-R (H-55) is a rabbit polyclonal antibody raised against amino acids 1-55 mapping at the N-terminus of MC5-R of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

MC5-R (H-55) is recommended for detection of MC5-R of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

MC5-R (H-55) is also recommended for detection of MC5-R in additional species, including equine, canine and porcine.

Suitable for use as control antibody for MC5-R siRNA (h): sc-35875, MC5-R shRNA Plasmid (h): sc-35875-SH and MC5-R shRNA (h) Lentiviral Particles: sc-35875-V.

MC5-R (H-55) X TransCruz antibody is recommended for ChIP assays.


Molecular Weight of MC5-R: 45 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200, K-562 whole cell lysate: sc-2203 or Hep G2 cell lysate: sc-2227.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



MC5-R (H-55): sc-28993. Western blot analysis of MC5-R expression in HeLa (A), K-562 (B), Hep G2 (C) and HL-60 (D) whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.