

# MC5-R (M-55): sc-28994

## BACKGROUND

The melanocortin family comprises the  $\alpha$ -,  $\beta$ - and  $\gamma$ -melanocyte stimulating hormones (MSH) and adrenocorticotrophin. The receptors for these hormones are seven-transmembrane, G protein-coupled proteins that activate adenyl cyclase. Five melanocortin receptors have been cloned and shown to exhibit different affinities and patterns of expression. MC1-R (MSH-R) is expressed in melanocytes and corticoadrenal tissue. MC2-R is the ACTH receptor and is expressed primarily in the adrenal cortex. MC3-R has been found in specific regions of the brain and is also expressed in placenta and gut. MC4-R is expressed primarily in brain, while MC5-R is expressed at low levels in most tissues.

## REFERENCES

1. Mountjoy, K.G., et al. 1992. The cloning of a family of genes that encode the melanocortin receptors. *Science* 257: 1248-1251.
2. Chhajlani, V., et al. 1992. Molecular cloning and expression of the human melanocyte stimulating hormone receptor cDNA. *FEBS Lett.* 309: 417-420.
3. Gantz, I., et al. 1993. Molecular cloning of a novel melanocortin receptor. *J. Biol. Chem.* 268: 8246-8250.
4. Gantz, I., et al. 1993. Molecular cloning, expression and gene localization of a fourth melanocortin receptor. *J. Biol. Chem.* 268: 15174-15179.
5. Roselli-Rehffuss, L., et al. 1993. Identification of a receptor for  $\gamma$ -melanotropin and other pro-opiomelanocortin peptides in the hypothalamus and limbic system. *Proc. Natl. Acad. Sci. USA* 90: 8856-8860.
6. Chhajlani, V., et al. 1993. Molecular cloning of a novel human melanocortin receptor. *Biochem. Biophys. Res. Commun.* 195: 866-873.
7. Mountjoy, K.G., et al. 1994. Localization of the melanocortin 4 receptor (MC4-R) in neuroendocrine and autonomic control circuits in the brain. *Mol. Endocrinol.* 8: 1298-1308.
8. Labbe, O., et al. 1994. Molecular cloning of a mouse melanocortin 5 receptor gene widely expressed in peripheral tissues. *Biochemistry* 33: 4543-4549.

## CHROMOSOMAL LOCATION

Genetic locus: MC5R (human) mapping to 18p11.2; Mc5r (mouse) mapping to 18 E2.

## SOURCE

MC5-R (M-55) is a rabbit polyclonal antibody raised against amino acids 1-55 mapping at the N-terminus of MC5-R of mouse origin.

## PRODUCT

Each vial contains 200  $\mu$ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## APPLICATIONS

MC5-R (M-55) is recommended for detection of MC5-R of mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1–2  $\mu$ g per 100–500  $\mu$ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


Suitable for use as control antibody for MC5-R siRNA (m): sc-35876.

Positive Controls: rat skeletal muscle extract.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/ 2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## DATA



MC5-R (M-55): sc-28994. Western blot analysis of mouse recombinant MC5-R fusion protein.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.