SANTA CRUZ BIOTECHNOLOGY, INC.

SRGAP1 (H-80): sc-292040



BACKGROUND

SRGAPs contain a highly conserved overall primary structure and play an important role in the cell facilitating Slit-robo signaling in cell migration and axon guidance. SRGAP1 (Slit-robo Rho GTPase activating protein 1), also known as ARHGAP13 (Rho GTPase activating protein 13), functions as a GTPase-activating protein for Cdc42 and Rho A. Expressed in kidney, testis, lung and brain, SRGAP1 contains an FCH (Fes/CIP4 homology) domain, a Rho-GAP domain and an SH3 domain. In the presence of Slit, SRGAP1 (via its SH3 domain) binds to the CC3 motif in robo (a protein responsible for mediating the repulsive effect of Slit) with higher affinity and inhibits Cdc42 activity in a robo/SRGAP-dependent manner. More specifically, SRGAP1 increases the intrinsic GTPase activity of Cdc42, thereby converting it to its inactive, GDP-bound form. Inactivation of Cdc42 ultimately leads to a decrease in actin polymerization.

REFERENCES

- Nagase, T., et al. 2000. Prediction of the coding sequences of unidentified human genes. XVI. The complete sequences of 150 new cDNA clones from brain which code for large proteins *in vitro*. DNA Res. 7: 65-73.
- Wong, K., et al. 2001. Signal transduction in neuronal migration: roles of GTPase activating proteins and the small GTPase Cdc42 in the Slit-Robo pathway. Cell 107: 209-221.
- 3. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 606523. World Wide Web URL: http://www.ncbi.nlm.nih.gov/omim/
- 4. Katoh, M., et al. 2003. FNBP2 gene on human chromosome 1q32.1 encodes ARHGAP family protein with FCH, FBH, Rho-GAP and SH3 domains. Int. J. Mol. Med. 11: 791-797.
- Katoh, M., et al. 2004. Identification and characterization of human FCH02 and mouse Fcho2 genes in silico. Int. J. Mol. Med. 14: 327-331.
- Katoh, Y., et al. 2004. Identification and characterization of ARHGAP27 gene in silico. Int. J. Mol. Med. 14: 943-947.
- 7. Katoh, M., et al. 2004. Characterization of human ARHGAP10 gene in silico. Int. J. Oncol. 25: 1201-1206.

CHROMOSOMAL LOCATION

Genetic locus: SRGAP1 (human) mapping to 12q14.2; Srgap1 (mouse) mapping to 10 D2.

SOURCE

SRGAP1 (H-80) is a rabbit polyclonal antibody raised against amino acids 831-910 mapping near the C-terminus of SRGAP1 of human origin.

PRODUCT

Each vial contains 200 μg lgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

SRGAP1 (H-80) is recommended for detection of SRGAP1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

SRGAP1 (H-80) is also recommended for detection of SRGAP1 in additional species, including equine, canine, bovine and porcine.


Suitable for use as control antibody for SRGAP1 siRNA (h): sc-95789, SRGAP1 siRNA (m): sc-153820, SRGAP1 shRNA Plasmid (h): sc-95789-SH, SRGAP1 shRNA Plasmid (m): sc-153820-SH, SRGAP1 shRNA (h) Lentiviral Particles: sc-95789-V and SRGAP1 shRNA (m) Lentiviral Particles: sc-153820-V.


Molecular Weight (predicted) of SRGAP1 isoforms: 124/122 kDa.

Molecular Weight (observed) of SRGAP1: 144-149 kDa.

Positive Controls: IMR-32 cell lysate: sc-2409, HEK293 whole cell lysate: sc-45136 or A549 cell lysate: sc-2413.

DATA





SRGAP1 expression in IMR-32 whole cell lysate

SRGAP1 (H-80): sc-292040. Western blot analysis of SRGAP1 expression in DU 145 $({\bf A}),$ HEK293 $({\bf B})$ and A549 $({\bf C})$ whole cell lysates.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS Satisfation Guaranteed

Try SRGAP1 (G-10): sc-390352 or SRGAP1 (D-11): sc-390349, our highly recommended monoclonal alternatives to SRGAP1 (H-80).