

Nop10 (H-54): sc-292092

BACKGROUND

Nop10, also known as NOLA3 (nucleolar protein family A member 3), is a 64 amino acid protein that localizes to the nucleolus, as well as to cajal bodies, and exists as a component of the multi-protein H/ACA small nucleolar ribonucleoprotein (H/ACA snoRNP) complex. Working in tandem with other members of the H/ACA snoRNP complex, Nop10 plays an essential role in telomere maintenance and ribosome biogenesis and is also thought to be required for the proper processing and trafficking of the TERT (telomerase reverse transcriptase) holoenzyme. Defects in the gene encoding Nop10 are the cause of dyskeratosis congenita autosomal recessive (ARDKC), a rare and progressive bone marrow failure syndrome that is characterized by reticulated skin hyperpigmentation, nail dystrophy and mucosal leukoplakia.

REFERENCES

- Henras, A., et al. 1998. Nhp2p and Nop10p are essential for the function of H/ACA snoRNPs. *EMBO J.* 17: 7078-7090.
- Pogaci, V., et al. 2000. Human H/ACA small nucleolar RNPs and telomerase share evolutionarily conserved proteins NHP2 and NOP10. *Mol. Cell. Biol.* 20: 9028-9040.
- Wang, C., et al. 2004. Architecture and assembly of mammalian H/ACA small nucleolar and telomerase ribonucleoproteins. *EMBO J.* 23: 1857-1867.
- Yamaguchi, H., et al. 2005. Mutations in TERT, the gene for telomerase reverse transcriptase, in aplastic anemia. *N. Engl. J. Med.* 352: 1413-1424.
- Hoareau-Aveilla, C., et al. 2006. hNaf1 is required for accumulation of human box H/ACA snoRNPs, scaRNPs, and telomerase. *RNA* 12: 832-840.
- Walne, A.J., et al. 2007. Genetic heterogeneity in autosomal recessive dyskeratosis congenita with one subtype due to mutations in the telomerase-associated protein NOP10. *Hum. Mol. Genet.* 16: 1619-1629.
- Online Mendelian Inheritance in Man, OMIM[™]. 2008. Johns Hopkins University, Baltimore, MD. MIM Number: 606471. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

CHROMOSOMAL LOCATION

Genetic locus: NOP10 (human) mapping to 15q14; Nop10 (mouse) mapping to 2 E3.

SOURCE

Nop10 (H-54) is a rabbit polyclonal antibody raised against amino acids 11-64 mapping at the C-terminus of Nop10 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Available as TransCruz reagent for Gel Supershift and ChIP applications, sc-292092 X, 200 µg/0.1 ml.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

Nop10 (H-54) is recommended for detection of Nop10 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Nop10 (H-54) is also recommended for detection of Nop10 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for Nop10 siRNA (h): sc-90082, Nop10 siRNA (m): sc-106308, Nop10 shRNA Plasmid (h): sc-90082-SH, Nop10 shRNA Plasmid (m): sc-106308-SH, Nop10 shRNA (h) Lentiviral Particles: sc-90082-V and Nop10 shRNA (m) Lentiviral Particles: sc-106308-V.

Nop10 (H-54) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.

Molecular Weight of Nop10: 8 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker[™] compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker[™] Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz[™] Mounting Medium: sc-24941.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **Nop10 (6H6): sc-517170**, our highly recommended monoclonal alternative to Nop10 (H-54).