

UVRAG (H-300): sc-292327

BACKGROUND

UVRAG (UV radiation resistance-associated gene), also known as p63 or DHTX, is a 699 amino acid cytoplasmic protein. UVRAG has been shown to activate the BECN1/PI 3-kinase complex, which promotes autophagy. Autophagy is the degradation of cellular proteins in the lysosomes and, when this pathway is suppressed, cell growth is deregulated. Mutations in the gene encoding UVRAG have been associated with colon cancer, suggesting that UVRAG is also involved in suppressing the proliferation and tumorigenicity of human colon cancer cells. UVRAG has been found to complement the ultraviolet sensitivity of xeroderma pigmentosum group C cells. Ubiquitously expressed, UVRAG is found at highest levels in kidney, lung, liver and brain. UVRAG contains one C2 domain, which is thought to be involved in calcium-dependent phospholipid binding.

CHROMOSOMAL LOCATION

Genetic locus: UVRAG (human) mapping to 11q13.5; Uvrage (mouse) mapping to 7 E2.

SOURCE

UVRAG (H-300) is a rabbit polyclonal antibody raised against amino acids 1-300 mapping at the N-terminus of UVRAG of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

UVRAG (H-300) is recommended for detection of UVRAG of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

UVRAG (H-300) is also recommended for detection of UVRAG in additional species, including equine, canine and bovine.

Suitable for use as control antibody for UVRAG siRNA (h): sc-76883, UVRAG siRNA (m): sc-76884, UVRAG shRNA Plasmid (h): sc-76883-SH, UVRAG shRNA Plasmid (m): sc-76884-SH, UVRAG shRNA (h) Lentiviral Particles: sc-76883-V and UVRAG shRNA (m) Lentiviral Particles: sc-76884-V.


Molecular Weight of UVRAG: 63 kDa.

Positive Controls: MCF7 whole cell lysate: sc-2206 or Ramos cell lysate: sc-2216.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



UVRAG (H-300): sc-292327. Western blot analysis of UVRAG expression in Ramos whole cell lysate.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.


 MONOS
Satisfaction
Guaranteed

Try **UVRAG (2E8): sc-293268**, our highly recommended monoclonal alternative to UVRAG (H-300).