



cathepsin D siRNA (h): sc-29239

BACKGROUND

The cathepsin family of proteolytic enzymes contains several diverse classes of proteases. The cysteine protease class comprises cathepsins B, L, H, K, S, and O. The aspartyl protease class is composed of cathepsins D and E. Cathepsin G is in the serine protease class. Most cathepsins are lysosomal and each is involved in cellular metabolism, participating in various events such as peptide biosynthesis and protein degradation. Cathepsins may also cleave some protein precursors, thereby releasing regulatory peptides. The promoter region of the cathepsin D gene contains five Sp1 binding sites and four AP-2 binding sites.

CHROMOSOMAL LOCATION

Genetic locus: CTSD (human) mapping to 11p15.5.

PRODUCT

cathepsin D siRNA (h) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see cathepsin D shRNA Plasmid (h): sc-29239-SH and cathepsin D shRNA (h) Lentiviral Particles: sc-29239-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

cathepsin D siRNA (h) is recommended for the inhibition of cathepsin D expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

cathepsin D (D-7): sc-377299 is recommended as a control antibody for monitoring of cathepsin D gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor cathepsin D gene expression knockdown using RT-PCR Primer: cathepsin D (h)-PR: sc-29239-PR (20 μ l, 439 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

SELECT PRODUCT CITATIONS

1. Sato, T., et al. 2008. Apoptosis supercedes necrosis in mitochondrial DNA-depleted Jurkat cells by cleavage of receptor-interacting protein and inhibition of lysosomal cathepsin. *J. Immunol.* 181: 197-207.
2. Byun, H.O., et al. 2009. Cathepsin D and eukaryotic translation elongation factor 1 as promising markers of cellular senescence. *Cancer Res.* 69: 4638-4647.
3. Liu, J.C., et al. 2009. Urotensin II induces rat cardiomyocyte hypertrophy via the transient oxidization of Src homology 2-containing tyrosine phosphatase and transactivation of epidermal growth factor receptor. *Mol. Pharmacol.* 76: 1186-1195.
4. Hu, S., et al. 2010. Prosaposin down-modulation decreases metastatic prostate cancer cell adhesion, migration, and invasion. *Mol. Cancer* 9: 30.
5. Chen, L., et al. 2011. Olfactomedin 4 suppresses prostate cancer cell growth and metastasis via negative interaction with cathepsin D and SDF-1. *Carcinogenesis* 32: 986-994.
6. Lin, C.F., et al. 2016. Glycogen synthase kinase-3 β and caspase-2 mediate ceramide- and etoposide-induced apoptosis by regulating the lysosomal-mitochondrial axis. *PLoS ONE* 11: e0145460.
7. Cocchiaro, P., et al. 2016. Lysosomal protease cathepsin D; a new driver of apoptosis during acute kidney injury. *Sci. Rep.* 6: 27112.
8. Huang, F., et al. 2019. Phosphorylation of nucleolin is indispensable to its involvement in the proliferation and migration of non-small cell lung cancer cells. *Oncol. Rep.* 41: 590-598.
9. Xie, Y., et al. 2021. Metformin and androgen receptor-axis-targeted (ARAT) agents induce two PARP-1-dependent cell death pathways in androgen-sensitive human prostate cancer cells. *Cancers* 13: 633.
10. Hsieh, Y.H., et al. 2021. Overexpression of lipocalin-2 inhibits proliferation and invasiveness of human glioblastoma multiforme cells by activating ERK targeting cathepsin D expression. *Biology* 10: 390.
11. Ashok, A., et al. 2021. Release of iron-loaded ferritin in sodium iodate-induced model of age related macular degeneration: an *in-vitro* and *in-vivo* study. *Antioxidants* 10: 1253.
12. Kitagawa, Y., et al. 2021. A novel topical fluorescent probe for detection of glioblastoma. *Clin. Cancer Res.* 27: 3936-3947.
13. Lee, S.G., et al. 2024. Cathepsin D promotes polarization of tumor-associated macrophages and metastasis through TGFBI-CCL20 signaling. *Exp. Mol. Med.* 56: 383-394.

RESEARCH USE

For research use only, not for use in diagnostic procedures.