

# Mig-2 (H-57): sc-292419

## BACKGROUND

Mig-2 recruits migfilin to cell-matrix adhesions, while the interaction with filamin mediates the association of migfilin with actin filaments. Together, Mig-2, migfilin and filamin define a connection between cell matrix adhesions and the actin cytoskeleton and participate in the orchestration of actin assembly and cell shape modulation. Mig-2 expression is transcriptionally elevated in leiomyomas and could be involved in its hormone-mediated growth of leiomyomas of the uterus. Expression of Mig-2 is ubiquitous, and it is found in numerous tumor tissues.

## REFERENCES

1. Stossel, T.P., et al. 2003. Filling gaps in signaling to Actin cytoskeletal remodeling. *Dev. Cell* 4: 444-445.
2. Tu Y., et al. 2003. Migfilin and Mig-2 link focal adhesions to filamin and the Actin cytoskeleton and function in cell shape modulation. *Cell* 113: 37-47.
3. Kato K., et al. 2004. Expression of the mitogen-inducible gene-2 (Mig-2) is elevated in human uterine leiomyomas but not in leiomyosarcomas. *Hum. Pathol.* 35: 55-60.
4. Tseng, Y., et al. 2004. The bimodal role of filamin in controlling the architecture and mechanics of F-Actin networks. *J. Biol. Chem.* 279: 1819-1826.
5. Pudas, R., et al. 2005. Structural basis for vertebrate filamin dimerization. *Structure* 13: 111-119.
6. Wu, C. 2005. Migfilin and its binding partners: from cell biology to human diseases. *J. Cell Sci.* 118: 659-664.
7. Gkretsi, V., et al. 2005. Physical and functional association of migfilin with cell-cell adhesions. *J. Cell Sci.* 118: 697-710.
8. SWISS-PROT/TrEMBL (Q96AC1). World Wide Web URL: <http://www.expasy.ch/sprot/sprot-top.html>

## CHROMOSOMAL LOCATION

Genetic locus: FERMT2 (human) mapping to 14q22.1; Fermt2 (mouse) mapping to 14 C1.

## SOURCE

Mig-2 (H-45) is a rabbit polyclonal antibody raised against amino acids 1-45 mapping at the N-terminus of Mig-2 of human origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.

## APPLICATIONS

Mig-2 (H-45) is recommended for detection of Mig-2 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000), may detect NP\_113659 fermitin family homolog 3.

Mig-2 (H-45) is also recommended for detection of Mig-2 in additional species, including equine, bovine and avian.

Suitable for use as control antibody for Mig-2 siRNA (h): sc-106786, Mig-2 siRNA (m): sc-149433, Mig-2 shRNA Plasmid (h): sc-106786-SH, Mig-2 shRNA Plasmid (m): sc-149433-SH, Mig-2 shRNA (h) Lentiviral Particles: sc-106786-V and Mig-2 shRNA (m) Lentiviral Particles: sc-149433-V.


Molecular Weight of Mig-2: 78 kDa.

Positive Controls: MDCK cell lysate: sc-2252 or PC-12 cell lysate: sc-2250.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## DATA



Mig-2 (H-57): sc-292419. Western blot analysis of Mig-2 expression in MDCK (A) and PC-12 (B) whole cell lysates.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

**MONOS**  
Satisfaction  
Guaranteed

Try **Mig-2 (NO-A16): sc-134387**, our highly recommended monoclonal alternative to Mig-2 (H-57).