

# NKG2-D (H-65): sc-292424

## BACKGROUND

The activity of natural killer (NK) cells is regulated by members of multiple receptor families that recognize class I MHC molecules, such as the killer cell inhibitory receptor/leukocyte immunoglobulin-like receptor (KIR/LIR) family and the C-type lectin superfamily. The KIR/LIR family includes p91A (also designated pp130 or PIR-B, for paired immunoglobulin-like receptor-B) and p91B (also designated PIR-A). p91A acts as an inhibitory receptor through interactions with SHP-1, whereas p91B acts as an activating receptor. CD94, NKG2 and Ly-49 are members of the C-type lectin superfamily of type II membrane glycoproteins. CD94 forms heterodimers with NKG2 isoforms on the surface of NK cells, whereas Ly-49 isoforms form homodimers. NKG2-D, expressed on NK cells,  $\gamma\delta$  T cells and CD8<sup>+</sup>  $\alpha\beta$  T cells, is a receptor for the stress inducible protein MICA, an antigen frequently expressed in epithelial tumors.

## REFERENCES

- Long, E.O. and Wagtmann, N. 1997. Natural killer cell receptors. *Curr. Opin. Immunol.* 9: 344-350.
- Moretta, A. and Moretta, L. 1997. HLA class I specific inhibitory receptors. *Curr. Opin. Immunol.* 9: 694-701.
- Hayami, K., Fukuta, D., Nishikawa, Y., Yamashita, Y., Inui, M., Ohyama, Y., Hikida, M., Ohmori, H. and Takai, T. 1997. Molecular cloning of a novel murine cell-surface glycoprotein homologous to killer cell inhibitory receptors. *J. Biol. Chem.* 272: 7320-7327.
- Ryan, J.C. and Seaman, W.E. 1997. Divergent functions of lectin-like receptors on NK cells. *Immunol. Rev.* 155: 79-89.
- Vance, R.E., Tanamachi, D.M., Hanke, T. and Raulet D.H. 1997. Cloning of a mouse homolog of CD94 extends the family of C-type lectins on murine natural killer cells. *Eur. J. Immunol.* 27: 3236-3241.
- Berg, K.L., Carlberg, K., Rohrschneider, L.R., Siminovitch, K.A. and Stanley, E.R. 1998. The major SHP-1-binding, tyrosine-phosphorylated protein in macrophages is a member of the KIR/LIR family and an SHP-1 substrate. *Oncogene* 17: 2535-2541.

## CHROMOSOMAL LOCATION

Genetic locus: KLRK1 (human) mapping to 12p13.2; Klrk1 (mouse) mapping to 6 F3.

## SOURCE

NKG2-D (H-65) is a rabbit polyclonal antibody raised against amino acids 108-172 mapping within an internal region of NKG2-D of human origin.

## PRODUCT

Each vial contains 200  $\mu$ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## APPLICATIONS

NKG2-D (H-65) is recommended for detection of NKG2-D of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for NKG2-D siRNA (h): sc-42948, NKG2-D siRNA (m): sc-42949, NKG2-D shRNA Plasmid (h): sc-42948-SH, NKG2-D shRNA Plasmid (m): sc-42949-SH, NKG2-D shRNA (h) Lentiviral Particles: sc-42948-V and NKG2-D shRNA (m) Lentiviral Particles: sc-42949-V.


Molecular Weight of NKG2-D: 42 kDa.

Positive Controls: K-562 whole cell lysate: sc-2203, WEHI-231 whole cell lysate: sc-2213 or TK-1 whole cell lysate: sc-364798.


## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## DATA



NKG2-D (H-65): sc-292424. Western blot analysis of NKG2-D expression in TK-1 (A) and WEHI-231 (B) whole cell lysates.



NKG2-D (H-65): sc-292424. Western blot analysis of NKG2-D expression in K-562 whole cell lysate.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.


 MONOS  
Satisfaction  
Guaranteed

Try **NKG2-D (5C6): sc-53501** or **NKG2-D (1D11): sc-23869**, our highly recommended monoclonal alternatives to NKG2-D (H-65).