

# UROD (H-53): sc-292502

## BACKGROUND

Uroporphyrinogen decarboxylase, also known as UROD or UPD, is a 367 amino acid protein that exists as a homodimer. UROD is the fifth enzyme in the human heme biosynthetic pathway and is responsible for the conversion of uroporphyrinogen to coproporphyrinogen through the removal of four carboxymethyl side chains. Mutations in the UROD gene are responsible for three autosomal disorders in humans: familial porphyria cutanea tarda (f-PCT), sporadic porphyria cutanea tarda (s-PCT) and hepatoerythropoietic porphyria (HEP). F-PCT is an autosomal dominant disorder characterized by late-onset light-sensitive dermatitis. High levels of uroporphyrin excretion in the urine and varying degrees of liver damage are associated with this disease. S-PCT is an idiosyncratic form of PCT that is characterized by a reduction of liver enzymes. HEP is an autosomal recessive disorder that affects infants. It is characterized by excessive excretion of acetate-substituted porphyrins and accumulation of protoporphyrin in erythrocytes.

## CHROMOSOMAL LOCATION

Genetic locus: UROD (human) mapping to 1p34.1; Urod (mouse) mapping to 4 D1.

## SOURCE

UROD (H-53) is a rabbit polyclonal antibody raised against amino acids 81-133 mapping within an internal region of UROD of human origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## APPLICATIONS

UROD (H-53) is recommended for detection of UROD of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

UROD (H-53) is also recommended for detection of UROD in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for UROD siRNA (h): sc-88548, UROD siRNA (m): sc-154937, UROD shRNA Plasmid (h): sc-88548-SH, UROD shRNA Plasmid (m): sc-154937-SH, UROD shRNA (h) Lentiviral Particles: sc-88548-V and UROD shRNA (m) Lentiviral Particles: sc-154937-V.


Molecular Weight of UROD: 41 kDa.

Positive Controls: K-562 whole cell lysate: sc-2203, Hep G2 cell lysate: sc-2227 or U-87 MG cell lysate: sc-2411.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## DATA



UROD (H-53): sc-292502. Western blot analysis of UROD expression in Hep G2 (A), K-562 (B), U-87 MG (C) and U-251 MG (D) whole cell lysates.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.


 MONOS  
Satisfaction  
Guaranteed

Try **UROD (G-7): sc-374318** or **UROD (C-4): sc-365297**, our highly recommended monoclonal alternatives to UROD (H-53).