

PFKL (H-36): sc-292523

BACKGROUND

Phosphofructokinases (PFKs) are regulatory glycolytic enzymes that catalyze the irreversible conversion of fructose-6-phosphate to fructose-1,6-bisphosphate. Mammalian PFK is a tetramer made up of diverse combinations of three isoenzymes: muscle (PFK-1), liver (PFKL) and platelet (PFKP). PFKL (phosphofructokinase, liver), also referred to as PFK-B (phosphofructo-1-kinase isozyme B), phosphofructokinase 1 or phosphohexokinase, predominates in organs with active gluconeogenesis, such as liver and kidney. Overexpression of PFKL in transgenic mice results in a diminished glucose-induced Insulin response, which suggests that PFKL may play a role in glucose-induced Insulin secretion. PFKL is expressed at high levels in Down's syndrome (DS) patients, suggesting a possible role for PFKL in the pathogenesis of DS.

REFERENCES

- Vora, S. and Francke, U. 1981. Assignment of the human gene for liver-type 6-phosphofructokinase isozyme (PFKL) to chromosome 21 by using somatic cell hybrids and monoclonal anti-L antibody. Proc. Natl. Acad. Sci. USA 78: 3738-3742.
- Levanon, D., et al. 1986. Genomic clones of the human liver-type phosphofructokinase. Biochem. Biophys. Res. Commun. 141: 374-380.
- Levanon, D., et al. 1987. Construction of a cDNA clone containing the entire coding region of the human liver-type phosphofructokinase. Biochem. Biophys. Res. Commun. 147: 1182-1187.
- Elson, A., et al. 1994. Overexpression of liver-type phosphofructokinase (PFKL) in transgenic-PFKL mice: implication for gene dosage in trisomy 21. Biochem. J. 299: 409-415.
- Knobler, H., et al. 1997. Impaired glucose-induced Insulin response in transgenic mice overexpressing the L-phosphofructokinase gene. Diabetes 46: 1414-1418.
- Peled-Kamar, M., et al. 1998. Altered brain glucose metabolism in transgenic-PFKL mice with elevated L-phosphofructokinase: *in vivo* NMR studies. Brain Res. 810: 138-145.

CHROMOSOMAL LOCATION

Genetic locus: PFKL (human) mapping to 21q22.3; Pfk1 (mouse) mapping to 10 C1.

SOURCE

PFKL (H-36) is a rabbit polyclonal antibody raised against amino acids 46-81 mapping near the N-terminus of PFKL of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

PFKL (H-36) is recommended for detection of PFKL of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000). PFKL (H-36) is also recommended for detection of PFKL in additional species, including equine, canine, bovine, porcine and avian.

Suitable for use as control antibody for PFKL siRNA (h): sc-106400, PFKL siRNA (m): sc-152180, PFKL shRNA Plasmid (h): sc-106400-SH, PFKL shRNA Plasmid (m): sc-152180-SH, PFKL shRNA (h) Lentiviral Particles: sc-106400-V and PFKL shRNA (m) Lentiviral Particles: sc-152180-V.


Molecular Weight of PFKL: 80 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200 or Hep G2 cell lysate: sc-2227.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



PFKL (H-36): sc-292523. Western blot analysis of PFKL expression in Hep G2 (A) and HeLa (B) whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

MONOS
Satisfaction
Guaranteed

Try **PFKL (A-6): sc-393713** or **PFKL (YT4): sc-100542**, our highly recommended monoclonal alternatives to PFKL (H-36).