

Rho GDI α (H-43): sc-292544

BACKGROUND

Members of the Ras superfamily of small GTP-binding proteins are critical mediators of diverse cell signaling pathways, including those leading to cell proliferation, cytoskeletal organization and secretion. The counter-conversion of the active GTP-bound form of these proteins to their inactive GDP-bound form is influenced by two types of regulatory proteins: those that alter the intrinsic GTPase activity of the GTP-binding proteins and those that alter the rate of GDP/GTP exchange. Guanine nucleotide-releasing factors (GRFs) increase the GDP dissociation rate, while GDP-dissociation inhibitors (GDIs) decrease the dissociation rate. Rho GDI α , also known as ARHGDI α or GDIA1, is a 204 amino acid member of the Rho GDI family of proteins. Localized to the cytoplasm, Rho GDI α inhibits the dissociation of GDP from Rho proteins, thereby preventing GTP from binding to and subsequently activating Rho proteins. In humans, Rho GDI α can be phosphorylated at Ser 101 by p21-activated kinase (α PAK), an event that inhibits Rho GDI α activity and may result in positive feedback regulation of certain Rho GDI α target proteins.

REFERENCES

1. Leffers, H., et al. 1993. Identification of two human Rho GDP dissociation inhibitor proteins whose overexpression leads to disruption of the Actin cytoskeleton. *Exp. Cell Res.* 209: 165-174.
2. Wagner, T., et al. 1997. A somatic cell hybrid panel for distal 17q: GDIA1 maps to 17q25.3. *Cytogenet. Cell Genet.* 76: 172-175.
3. Di-Poi, N., et al. 2001. Mechanism of NADPH oxidase activation by the Rac/Rho GDI complex. *Biochemistry* 40: 10014-10022.
4. Online Mendelian Inheritance in Man, OMIM[™]. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 601925. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
5. DerMardirossian, C., et al. 2004. Phosphorylation of Rho GDI by Pak1 mediates dissociation of Rac GTPase. *Mol. Cell* 15: 117-127.
6. Dransart, E., et al. 2005. Uncoupling of inhibitory and shuttling functions of rho GDP dissociation inhibitors. *J. Biol. Chem.* 280: 4674-4683.
7. DerMardirossian, C., et al. 2006. Phosphorylation of Rho GDI by Src regulates Rho GTPase binding and cytosol-membrane cycling. *Mol. Biol. Cell* 17: 4760-4768.
8. El Marzouk, S., et al. 2007. Rho GDP dissociation inhibitor α interacts with estrogen receptor α and influences estrogen responsiveness. *J. Mol. Endocrinol.* 39: 249-259.

CHROMOSOMAL LOCATION

Genetic locus: ARHGDI α (human) mapping to 17q25.3; Arhgdia (mouse) mapping to 11 E2.

SOURCE

Rho GDI α (H-43) is a rabbit polyclonal antibody raised against amino acids 2-44 mapping at the N-terminus of Rho GDI α of human origin.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

Rho GDI α (H-43) is recommended for detection of Rho GDI α of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


Rho GDI α (H-43) is also recommended for detection of Rho GDI α in additional species, including equine, canine, bovine, porcine and avian.

Suitable for use as control antibody for Rho GDI α siRNA (h): sc-36417, Rho GDI α siRNA (m): sc-36416, Rho GDI α siRNA (r): sc-61880, Rho GDI α shRNA Plasmid (h): sc-36417-SH, Rho GDI α shRNA Plasmid (m): sc-36416-SH, Rho GDI α shRNA Plasmid (r): sc-61880-SH, Rho GDI α shRNA (h) Lentiviral Particles: sc-36417-V, Rho GDI α shRNA (m) Lentiviral Particles: sc-36416-V and Rho GDI α shRNA (r) Lentiviral Particles: sc-61880-V.

Molecular Weight of Rho GDI α : 30 kDa.

Positive Controls: PC-12 cell lysate: sc-2250, HeLa whole cell lysate: sc-2200 or SK-BR-3 cell lysate: sc-2218.

DATA



Rho GDI α (H-43): sc-292544. Western blot analysis of Rho GDI α expression in PC-12 (A) and SK-BR-3 (B) whole cell lysates.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try Rho GDI α (G-2): sc-373724 or Rho GDI α (B-10): sc-13120, our highly recommended monoclonal alternatives to Rho GDI α (H-43).