

# AP-4 (H-84): sc-292568

## BACKGROUND

AP-2 transcription factor family members include AP-2 $\alpha$ , AP-2 $\beta$  and AP-2 $\gamma$ , which specifically bind to the DNA consensus sequence CCCCAGGC and initiate transcription of selected genes. AP-2, also known as ERF-1, plays a role in regulating estrogen receptor expression. AP-2 $\beta$ , a splice variant of AP-2 $\alpha$ , inhibits AP-2 activity. Besides subscribing to the AP-2 complex, AP-2 $\alpha$ , AP-2 $\beta$  and AP-2 $\gamma$  proteins compose the OB2-1 transcription factor complex. OB2-1 specifically upregulates expression of the proto-oncogene c-ErbB-2, which is overexpressed in 25-30% of breast cancers. The gene encoding AP-2 $\alpha$  maps to human chromosome 6p24.3. AP-2 $\alpha$  may play an important role in the development of ectodermal-derived tissues. Deleterious mutations involving the AP-2 $\alpha$  gene are linked to microphthalmia, corneal clouding and other anterior eye chamber defects. The ubiquitously expressed AP-4 transcription factor specifically binds to the DNA consensus sequence 5'-CAGCTG-3'. AP-4 interacts with promoters for immunoglobulin- $\kappa$  gene families and simian virus 40. AP-4 may enhance the transcription of the human Huntington's disease gene. AP-4 is a helix-loop-helix protein that contains two distinctive leucine repeat elements.

## CHROMOSOMAL LOCATION

Genetic locus: TFAP4 (human) mapping to 16p13.3; Tcfap4 (mouse) mapping to 16 A1.

## SOURCE

AP-4 (H-84) is a rabbit polyclonal antibody raised against amino acids 93-176 mapping within an internal region of AP-4 of human origin.

## PRODUCT

Each vial contains 200  $\mu$ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## APPLICATIONS

AP-4 (H-84) is recommended for detection of AP-4 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

AP-4 (H-84) is also recommended for detection of AP-4 in additional species, including equine, canine and bovine.

Suitable for use as control antibody for AP-4 siRNA (h): sc-37690, AP-4 siRNA (m): sc-37691, AP-4 shRNA Plasmid (h): sc-37690-SH, AP-4 shRNA Plasmid (m): sc-37691-SH, AP-4 shRNA (h) Lentiviral Particles: sc-37690-V and AP-4 shRNA (m) Lentiviral Particles: sc-37691-V.


Molecular Weight of AP-4: 48 kDa.

Positive Controls: IMR-32 nuclear extract: sc-2148 or Jurkat nuclear extract: sc-2132.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## DATA



AP-4 (H-84): sc-292568. Western blot analysis of AP-4 expression in IMR-32 (A) and Jurkat (B) nuclear extracts.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.

**MONOS**  
Satisfaction  
Guaranteed

Try **AP-4 (A-8): sc-377042** or **AP-4 (F-1): sc-376977**, our highly recommended monoclonal alternatives to AP-4 (H-84).