

ERp72 (H-272): sc-292586

BACKGROUND

Mammals defend themselves against intracellular pathogens through presentation of cytoplasmically derived short pathogenic peptides to the cell surface of cytotoxic T lymphocytes, which subsequently leads to cytotoxic events with respect to the affected cell. Antigen presentation is mediated by major histocompatibility complex (MHC) class I molecules, which bind and coordinate short pathogenic peptides. The proper folding and assembly of MHC class I molecules in the endoplasmic reticulum (ER) involve a number of components. MHC class I molecules assemble in the ER with chaperones before binding to the transporter associated with antigen processing (TAP) protein. ERp57 is a component of the MHC class I pathway that appears to interact with MHC class I molecules before they associate with TAP. ERp72, also designated protein disulfide-isomerase A4, is involved in the catalysis of protein-S-S- bond rearrangement. ERp57 and ERp72 may act as proteases, protein disulfide isomerases, phospholipases or a combination of these.

CHROMOSOMAL LOCATION

Genetic locus: PDIA4 (human) mapping to 7q36.1; Pdia4 (mouse) mapping to 6 B2.3.

SOURCE

ERp72 (H-272) is a rabbit polyclonal antibody raised against amino acids 257-528 mapping within an internal region of ERp72 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

ERp72 (H-272) is recommended for detection of ERp72 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

ERp72 (H-272) is also recommended for detection of ERp72 in additional species, including porcine.

Suitable for use as control antibody for ERp72 siRNA (h): sc-44571, ERp72 siRNA (m): sc-44576, ERp72 shRNA Plasmid (h): sc-44571-SH, ERp72 shRNA Plasmid (m): sc-44576-SH, ERp72 shRNA (h) Lentiviral Particles: sc-44571-V and ERp72 shRNA (m) Lentiviral Particles: sc-44576-V.


Molecular Weight of ERp72: 72 kDa.

Positive Controls: Jurkat whole cell lysate: sc-2204, Caki-1 cell lysate: sc-2224 or JAR cell lysate: sc-2276.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 4) Immunohistochemistry: use ImmunoCruz™: sc-2051 or ABC: sc-2018 rabbit IgG Staining Systems.

DATA



ERp72 (H-272): sc-292586. Western blot analysis of ERp72 expression in Jurkat (A), Caki-1 (B), JAR (C), HeLa (D), K-562 (E) and Ramos (F) whole cell lysates.



ERp72 (H-272): sc-292586. Immunoperoxidase staining of formalin fixed, paraffin-embedded human liver tissue showing cytoplasmic staining of hepatocytes.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS
Satisfaction
Guaranteed

Try **ERp72 (B-4): sc-390530** or **ERp72 (C-7): sc-376230**, our highly recommended monoclonal alternatives to ERp72 (H-272).