

# PUS1 (H-280): sc-292591

## BACKGROUND

PUS1 (pseudouridine synthase 1) belongs to the tRNA pseudouridine synthase truA family. PUS1 functions in the conversion of uridine into pseudouridine after the nucleotide has been incorporated into RNA. It may have a functional role in tRNAs and is also thought to assist in the peptidyl transfer reaction of rRNAs. As a nucleus-resident protein, PUS1 forms a complex with RARG and the SRA1 RNA. PUS1 is widely expressed, with highest levels of expression in the brain and skeletal muscle tissues. Defects in PUS1 are a cause of myopathy with lactic acidosis and sideroblastic anemia (MLASA), also known as mitochondrial myopathy and sideroblastic anemia. MLASA is a rare autosomal recessive oxidative phosphorylation disorder specific to bone marrow and skeletal muscle. The deduced human PUS1 protein contains 348 amino acids and shares 92% sequence homology with mouse PUS1.

## CHROMOSOMAL LOCATION

Genetic locus: PUS1 (human) mapping to 12q24.33; Pus1 (mouse) mapping to 5 F.

## SOURCE

PUS1 (H-280) is a rabbit polyclonal antibody raised against amino acids 148-427 mapping at the C-terminus of PUS1 of human origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## APPLICATIONS

PUS1 (H-280) is recommended for detection of PUS1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

PUS1 (H-280) is also recommended for detection of PUS1 in additional species, including equine, bovine and porcine.

Suitable for use as control antibody for PUS1 siRNA (h): sc-61417, PUS1 siRNA (m): sc-61418, PUS1 shRNA Plasmid (h): sc-61417-SH, PUS1 shRNA Plasmid (m): sc-61418-SH, PUS1 shRNA (h) Lentiviral Particles: sc-61417-V and PUS1 shRNA (m) Lentiviral Particles: sc-61418-V.


Molecular Weight of PUS1 isoform 1/2: 47/44 kDa.

Positive Controls: A-431 whole cell lysate: sc-2201, NCI-H460 whole cell lysate: sc-364235 or Jurkat whole cell lysate: sc-2204.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## DATA



PUS1 (H-280): sc-292591. Western blot analysis of PUS1 expression in A-431 (A), NCI-H460 (B) and Jurkat (C) whole cell lysates.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.


 MONOS  
Satisfaction  
Guaranteed

Try **PUS1 (A-4): sc-390043** or **PUS1 (B-6): sc-376443**, our highly recommended monoclonal alternatives to PUS1 (H-280).