# MFAP3L (H-164): sc-292672



The Power to Question

## **BACKGROUND**

MFAP3L (microfibrillar-associated protein 3-like), also known as HSD39 or testis development protein NYD-SP9, is a 409 amino acid single-pass type I cell membrane protein that contains one Ig-like (immunoglobulin-like) domain. Found primarily in testis, MFAP3L is encoded by a gene that is located on chromosome 4 and is expressed as three isoforms due to alternative splicing events. Representing approximately 6% of the human genome, chromosome 4 contains nearly 900 genes, one of which is the Huntingtin gene, which is found to encode an expanded glutamine tract in cases of Huntington's disease. FGFR-3 is also encoded on chromosome 4 and has been associated with thanatophoric dwarfism, achondroplasia, Muenke syndrome and bladder cancer. Chromosome 4 is also tied to Ellis-van Creveld syndrome, methylmalonic acidemia and polycystic kidney disease.

# **REFERENCES**

- Hillier, L.W., et al. 2005. Generation and annotation of the DNA sequences of human chromosomes 2 and 4. Nature 434: 724-731.
- 2. Cowan, C.M. and Raymond, L.A. 2006. Selective neuronal degeneration in Huntington's disease. Curr. Top. Dev. Biol. 75: 25-71.
- Chandler, R.J., et al. 2007. Metabolic phenotype of methylmalonic acidemia in mice and humans: the role of skeletal muscle. BMC Med. Genet. 8: 64.
- Cunningham, M.L., et al. 2007. Syndromic craniosynostosis: from history to hydrogen bonds. Orthod. Craniofac. Res. 10: 67-81.
- Doherty, E.S., et al. 2007. Muenke syndrome (FGFR3-related craniosynostosis): Expansion of the phenotype and review of the literature. Am. J. Med. Genet. A 143: 3204-3215.
- Ruiz-Perez, V.L., et al. 2007. Evc is a positive mediator of lhh-regulated bone growth that localises at the base of chondrocyte cilia. Development 134: 2903-2912.
- 7. Stack, E.C., et al. 2007. Neuroprotective effects of synaptic modulation in Huntington's disease R6/2 mice. J. Neurosci. 27: 12908-12915.
- 8. Versteegh, F.G., et al. 2007. Growth hormone analysis and treatment in Ellis-van Creveld syndrome. Am. J. Med. Genet. Part A 143A: 2113-2121.

## CHROMOSOMAL LOCATION

Genetic locus: MFAP3L (human) mapping to 4q33; Mfap3l (mouse) mapping to 8 B3.1.

## SOURCE

MFAP3L (H-164) is a rabbit polyclonal antibody raised against amino acids 246-409 mapping at the C-terminus of MFAP3L of human origin.

# **PRODUCT**

Each vial contains 200  $\mu g$  lgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

# **RESEARCH USE**

For research use only, not for use in diagnostic procedures.

#### **APPLICATIONS**

MFAP3L (H-164) is recommended for detection of MFAP3L isoforms 1 and 2 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

MFAP3L (H-164) is also recommended for detection of MFAP3L isoforms 1 and 2 in additional species, including equine and canine.

Suitable for use as control antibody for MFAP3L siRNA (h): sc-88992, MFAP3L siRNA (m): sc-149402, MFAP3L shRNA Plasmid (h): sc-88992-SH, MFAP3L shRNA Plasmid (m): sc-149402-SH, MFAP3L shRNA (h) Lentiviral Particles: sc-88992-V and MFAP3L shRNA (m) Lentiviral Particles: sc-149402-V.

Molecular Weight of MFAP3L: 45 kDa.

Positive Controls: Jurkat whole cell lysate: sc-2204.

## **RECOMMENDED SECONDARY REAGENTS**

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

#### **STORAGE**

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

# **PROTOCOLS**

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

Santa Cruz Biotechnology, Inc. 1.800.457.3801 831.457.3801 Fax 831.457.3801 Europe +00800 4573 8000 49 6221 4503 0 www.scbt.com