# SANTA CRUZ BIOTECHNOLOGY, INC.

# goat anti-chicken IgY-AP: sc-2928



### BACKGROUND

Santa Cruz Biotechnology's secondary antibodies are available conjugated to either an enzyme, biotin or fluorophore for use in a variety of antibody-based applications including Western Blot, immunostaining, flow cytometry and ELISA. Secondary antibodies are commonly affinity purified against immobilized whole IgG isotypes, including IgG<sub>1</sub>, IgG<sub>2a</sub>, IgG<sub>2b</sub>, IgG<sub>3</sub> and IgG<sub>4</sub>. Santa Cruz Biotechnology offers a wide selection of secondary antibodies, which are used in conjunction with our Cruz Marker<sup>TM</sup> molecular weight standards. We also provide specialized secondaries, such as pre-adsorbed secondary antibodies, which are pre-adsorbed with human IgG and mouse IgG for immuno-globulin-rich tissues and cells,  $F(ab')_2$  fragment secondary antibodies that reduce non-specific secondary antibody binding to Fc receptors on the cell surface, and isotype-specific secondary antibodies against IgM, IgA and IgY.

## SOURCE

goat anti-chicken IgY-AP is a pre-adsorbed, affinity purified secondary antibody raised in goat against chicken IgY and conjugated to AP (alkaline phosphatase).

## PRODUCT

Each vial contains 200  $\mu g$  mouse and human adsorbed IgY in 0.5 ml of PBS containing 50% glycerol, 1 mM zinc chloride, 0.02% sodium azide and 1 mM magnesium chloride.

### **APPLICATIONS**

goat anti-chicken IgY-AP is recommended for detection of chicken IgY by Western Blotting of immunoglobulin-rich tissues and cells (starting dilution: 1:5000, dilution range 1:5000-1:10000).

# **RECOMMENDED SUPPORT PRODUCTS**

- <sup>n</sup> Western Blotting Luminol Reagent, for 2,000 cm<sup>2</sup> membrane area: sc-2048
- n RIPA Lysis Buffer, 50 ml, cell lysis buffer with protease inhibitors: sc-24948
- Electrophoresis Sample Buffer, 2X, 25 ml, reducing buffer: sc-24945
- <sup>n</sup> Complete<sup>™</sup> Protease Inhibitor Cocktail Tablet, 20 tablets: sc-29130
- Running Buffer, 10X, 1 L, TRIS-Glycine WB running buffer, pH 8.3: sc-24949
- <sup>n</sup> Towbin, with SDS, 10X, 1 L, WB transfer buffer pH 8.3: sc-24954
- Bovine Serum Albumin (BSA), 100 g, blocking/incubation agent: sc-2323
- <sup>n</sup> TBS Blotto A, lyophilized powder in single-use bottle: sc-2333
- <sup>¬</sup> UltraCruz™ PVDF Transfer Membrane, 0.45 μm, 30 cm x 3 m roll: sc-3723
- □ UltraCruz<sup>™</sup> Nitrocellulose Pure Transfer Membrane, 0.22 µm, 30 cm x 3 m roll: sc-3718
- <sup>n</sup> UltraCruz™ Tissue Culture Dish, 100 mm polystyrene dish: sc-200286
- <sup>n</sup> UltraCruz<sup>™</sup> Cell Scrapers, 25 cm, sterile, 100 per case: sc-213229
- □ UltraCruz™ Electrophoresis Cell: sc-201625: runs up to 10 or 15 sample by SDS – PAGE protein electrophoresis
- <sup>\_</sup> UltraCruz<sup>™</sup> Autoradiography Film, Blue, 8 x 1, 100 sheets: sc-201697
- □ UltraCruz™ Gel Incubation Trays, 100 per pack: sc-201755 (blue), sc-201756 (green), sc-201757 (pink), sc-201758 (yellow), sc-201759 (orange)

## SELECT PRODUCT CITATIONS

 Yalvac, M.E., et al. 2011. Differentiation and neuro-protective properties of immortalized human tooth germ stem cells. Neurochem. Res. 36: 2227-2235.

## **STORAGE**

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## **RESEARCH USE**

For research use only, not for use in diagnostic procedures.

#### **PROTOCOLS**

See our web site at www.scbt.com or our catalog for detailed protocols and support products.