

Tak1L (H-214): sc-292803

BACKGROUND

Several serine/threonine protein kinases have been implicated as intermediates in signal transduction pathways. These include ERK/MAP kinases, ribosomal S6 kinase (Rsk) and Raf-1. Raf-1 is a protein with intrinsic kinase activity towards serine/threonine residues and is widely expressed in many tissue types and cell lines. Raf-1 activation is dependent on the small molecular weight GTPase Ras. Two proteins putatively involved in this process are Ksr-1 and Tak1. Ksr-1 (kinase suppressor of Ras) is a novel Raf-related protein kinase whose function is required for Ras signal transduction. Whether Ksr-1 lies directly downstream of Ras or acts in a parallel pathway is not yet known. Tak1 (TGF β -activated kinase) has been shown to participate in the activation of the MAP kinase family in response to TGF β stimulation. Tak1L (Tak1-like protein), also known as C21orf7, is a 242 amino acid protein that shares homology with the C-terminal tail of Tak1. Tak1L is expressed predominantly in peripheral blood leukocytes, with strong expression found in the adenocarcinomic cell lines GI-112 and PC-3 and the carcinomic cell line GI-101.

REFERENCES

1. Huleihel, M., et al. 1986. Characterization of murine A-Raf, a new oncogene related to the v-Raf oncogene. *Mol. Cell. Biol.* 6: 2655-2662.
2. Ray, L.B. and Sturgill, T.W. 1988. Insulin-stimulated microtubule-associated protein kinase is phosphorylated on tyrosine and threonine *in vivo*. *Proc. Natl. Acad. Sci. USA* 85: 3753-3757.
3. Morrison, D.K., et al. 1988. Signal transduction from membrane to cytoplasm: growth factors and membrane-bound oncogene products increase Raf-1 phosphorylation and associated protein kinase activity. *Proc. Natl. Acad. Sci. USA* 85: 8855-8859.
4. Pelech, S.L., et al. 1990. Protein kinase cascades in meiotic and mitotic cell cycle control. *Biochem. Cell Biol.* 68: 1297-1330.
5. Downward, J. 1995. Ksr: a novel player in the Ras pathway. *Cell* 83: 831-834.

CHROMOSOMAL LOCATION

Genetic locus: C21orf7 (human) mapping to 21q21.3; ORF63 (mouse) mapping to 16 C3.3.

SOURCE

Tak1L (H-214) is a rabbit polyclonal antibody raised against amino acids 29-242 mapping at the C-terminus of Tak1L of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

Tak1L (H-214) is recommended for detection of Tak1L of human and, to a lesser extent, mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Tak1L (H-214) is also recommended for detection of Tak1L in additional species, including equine.

Suitable for use as control antibody for Tak1L siRNA (h): sc-91477, Tak1L siRNA (m): sc-154063, Tak1L shRNA Plasmid (h): sc-91477-SH, Tak1L shRNA Plasmid (m): sc-154063-SH, Tak1L shRNA (h) Lentiviral Particles: sc-91477-V and Tak1L shRNA (m) Lentiviral Particles: sc-154063-V.

Molecular Weight (predicted) of Tak1L isoforms: 27/15/16 kDa.

Molecular Weight (observed) of human Tak1L: 30 kDa.

Molecular Weight (observed) of mouse Tak1L: 32/22 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker[™] compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker[™] Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz[™] Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.