

M-Ras (H-60): sc-292844

BACKGROUND

The mammalian c-H-, c-K- and N-Ras proto-oncogenes encode proteins that are ubiquitously expressed in vertebrate cells. c-H- and c-K-Ras are cellular homologs of the v-H- and v-K-Ras sequences originally isolated from the Harvey and Kirsten strains of rat sarcoma virus. p21Ras-encoded proteins bind GDP and GTP with high affinity and possess a low level intrinsic GTPase activity that can be stimulated over 100-fold by interaction with cytosolic GTPase activating protein (GAP), a potential effector for Ras p21 function. Point mutations at amino acids 12, 13, 59 and 61 within domains responsible for GTP binding and hydrolysis activate Ras proteins to their oncogenic form and block the ability of their GTPase activities to be stimulated by GAP. M-Ras has been identified as a GTPase that shares structural similarities to the Ras family proteins. M-Ras is thought to play a role in reorganization of the actin cytoskeleton.

REFERENCES

- Shih, T.Y., Papageorge, A.G., Stokes, P.E., Weeks, M.O. and Scolnick, E.M. 1980. Guanine nucleotide-binding and autophosphorylating activities associated with the p21src protein of Harvey murine sarcoma virus. *Nature* 287: 686-691.
- Ellis, R.W., DeFeo, D., Shih, T.Y., Gonda, M.A., Young, H.A., Tsuchida, N., Lowy, D.R. and Scolnick, E.M. 1981. The p21 src genes of Harvey and Kirsten sarcoma viruses originate from divergent members of a family of normal vertebrate genes. *Nature* 292: 506-511.
- Barbacid, M. 1987. ras genes. *Ann. Rev. Biochem.* 56: 779-827.
- Trahey, M. and McCormick, F. 1987. A cytoplasmic protein stimulates normal N-ras p21 GTPase, but does not affect oncogenic mutants. *Science* 238: 542-545.
- Cales, C., Hancock, J.F., Marshall, C.J. and Hall, A. 1988. The cytoplasmic protein GAP is implicated as the target for regulation by the ras gene product. *Nature* 332: 548-551.
- Adari, H., Lowy, D., Willumsen, B.M., Der, C.J. and McCormick, F. 1988. Guanosine triphosphatase activating protein (GAP) interacts with the p21 ras effector binding domain. *Science* 240: 518-521.
- Matsumoto, K., Asano, T. and Endo, T. 1997. Novel small GTPase M-Ras participates in reorganization of actin cytoskeleton. *Oncogene* 15: 2409-2417.

CHROMOSOMAL LOCATION

Genetic locus: MRAS (human) mapping to 3q22.3; Mras (mouse) mapping to 9 E3.3.

SOURCE

M-Ras (H-60) is a rabbit polyclonal antibody raised against amino acids 114-173 mapping near the C-terminus of M-Ras of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

M-Ras (H-60) is recommended for detection of M-Ras of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

M-Ras (H-60) is also recommended for detection of M-Ras in additional species, including equine, canine, bovine, porcine and avian.

Suitable for use as control antibody for M-Ras siRNA (h): sc-41857, M-Ras siRNA (m): sc-41858, M-Ras shRNA Plasmid (h): sc-41857-SH, M-Ras shRNA Plasmid (m): sc-41858-SH, M-Ras shRNA (h) Lentiviral Particles: sc-41857-V and M-Ras shRNA (m) Lentiviral Particles: sc-41858-V.


Molecular Weight of M-Ras: 29 kDa.

Positive Controls: Jurkat whole cell lysate: sc-2204.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



M-Ras (H-60): sc-292844. Western blot analysis of M-Ras expression in Jurkat whole cell lysate.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.