

# dNT-1 (H-48): sc-292986

## BACKGROUND

Deoxyribonucleotidases are catabolic proteins that regulate intracellular deoxyribonucleoside triphosphate pools through substrate cycles. The various substrate specificities of deoxyribonucleotidases suggests that these enzymes have different functions in nucleotide metabolism. For example, dNT-2 is a mitochondrial specific enzyme that regulates a thymidine/dTMP substrate cycle by catalyzing the dephosphorylation of 5'- and 2'(3')-phosphates of uracil and thymine, thereby regulating the size of the intramitochondrial dTTP pool. Human dNT-1 is a cytosolic enzyme that regulates pyrimidine nucleotide pools. Human dNT-2 contains a mitochondrial leader peptide, providing the structural basis for two-step processing during import into the mitochondrial matrix. Mitochondrial dNT-2 is 52% identical to cytosolic deoxyribonucleotidase (dNT-1) and the two enzymes share many catalytic properties, however dNT-2 shows a more narrow substrate specificity. The human dNT-2 gene maps to chromosome 17p11.2, which is also a critical region for the Smith-Magenis syndrome, suggesting that dNT-2 may be involved in the etiology of this hereditary disease.

## REFERENCES

- Rampazzo, C., Johansson, M., Gallinaro, L., Ferraro, P., Hellman, U., Karlsson, A., Reichard, P. and Bianchi, V. 2000. Mammalian 5'(3')-deoxyribonucleotidase, cDNA cloning, and overexpression of the enzyme in *Escherichia coli* and mammalian cells. *J. Biol. Chem.* 275: 5409-5415.
- Rampazzo, C., Gallinaro, L., Milanese, E., Frigimelica, E., Reichard, P. and Bianchi, V. 2000. A deoxyribonucleotidase in mitochondria: involvement in regulation of dNTP pools and possible link to genetic disease. *Proc. Natl. Acad. Sci. USA* 97: 8239-8244.
- Gazziola, C., Ferraro, P., Moras, M., Reichard, P. and Bianchi, V. 2001. Cytosolic high K(m) 5'-nucleotidase and 5'(3')-deoxyribonucleotidase in substrate cycles involved in nucleotide metabolism. *J. Biol. Chem.* 276: 6185-6190.
- Online Mendelian Inheritance in Man, OMIM™. 2001. Johns Hopkins University, Baltimore, MD. MIM Number: 605292. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- LocusLink Report (LocusID: 56953). <http://www.ncbi.nlm.nih.gov/LocusLink/>

## CHROMOSOMAL LOCATION

Genetic locus: NT5C (human) mapping to 17q25.1; Nt5c (mouse) mapping to 11 E2.

## SOURCE

dNT-1 (H-48) is a rabbit polyclonal antibody raised against amino acids 129-176 mapping within an internal region of dNT-1 of human origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## APPLICATIONS

dNT-1 (H-48) is recommended for detection of dNT-1 of mouse and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


dNT-1 (H-48) is also recommended for detection of dNT-1 in additional species, including equine, canine and porcine.

Suitable for use as control antibody for dNT-1 siRNA (h): sc-38995, dNT-1 siRNA (m): sc-38996, dNT-1 shRNA Plasmid (h): sc-38995-SH, dNT-1 shRNA Plasmid (m): sc-38996-SH, dNT-1 shRNA (h) Lentiviral Particles: sc-38995-V and dNT-1 shRNA (m) Lentiviral Particles: sc-38996-V.


Molecular Weight of dNT-1 isoforms: 23/13 kDa.

Positive Controls: dNT-1 (h): 293 Lysate: sc-110915, Raji whole cell lysate: sc-364236 or RT-4 whole cell lysate: sc-364257.

## DATA



dNT-1 (H-48): sc-292986. Western blot analysis of dNT-1 expression in Raji (A), RT-4 (B) and U-251-MG (C) whole cell lysates and human liver tissue extract (D).



dNT-1 (H-48): sc-292986. Western blot analysis of dNT-1 expression in non-transfected 293: sc-110760 (A), human dNT-1 transfected 293: sc-110915 (B) and MIA PaCa-2 (C) whole cell lysates.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.



Try **dNT-1 (C-10): sc-390041** or **dNT-1 (F-9): sc-271972**, our highly recommended monoclonal alternatives to dNT-1 (H-48).