

PBK (11): sc-293028

BACKGROUND

Protein kinases comprise a large group of encoded factors that regulate cellular processes by catalyzing the transfer of a phosphate group to a hydroxyl acceptor in serine, threonine or tyrosine residues. Kinases are capable of influencing the oncogenic potential of cell systems at the level of oncoprotein or tumor suppressor protein phosphorylation states. Human PDZ-binding kinase, known as PBK, is a 322 amino acid, T/SXV motif-containing serine/threonine kinase that is abundant in placenta and absent from adult brain tissue. A PDZ domain in the tumor suppressor protein Dlg can coordinate with the T/SXV motif of PBK. The cell cycle checkpoint kinase Cdc2/cyclin B is an upstream effector of PBK that can phosphorylate and activate PBK. Active PBK may associate with PDZ-containing proteins and influence cell cycle control or cellular proliferation.

CHROMOSOMAL LOCATION

Genetic locus: PBK (human) mapping to 8p21.1; Pbk (mouse) mapping to 14 D1.

SOURCE

PBK (11) is a mouse monoclonal antibody raised against amino acids 37-252 of PBK of mouse origin.

PRODUCT

Each vial contains 200 µg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

PBK (11) is available conjugated to agarose (sc-293028 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-293028 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-293028 PE), fluorescein (sc-293028 FITC), Alexa Fluor® 488 (sc-293028 AF488), Alexa Fluor® 594 (sc-293028 AF594) or Alexa Fluor® 647 (sc-293028 AF647), 200 µg/ml, for IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-293028 AF680) or Alexa Fluor® 790 (sc-293028 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

APPLICATIONS

PBK (11) is recommended for detection of PBK of mouse, rat, human and canine origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500).

Suitable for use as control antibody for PBK siRNA (h): sc-106892, PBK siRNA (m): sc-152042, PBK shRNA Plasmid (h): sc-106892-SH, PBK shRNA Plasmid (m): sc-152042-SH, PBK shRNA (h) Lentiviral Particles: sc-106892-V and PBK shRNA (m) Lentiviral Particles: sc-152042-V.

Molecular Weight of PBK: 36 kDa.

Positive Controls: 3T3-L1 cell lysate: sc-2243, F9 cell lysate: sc-2245 or RAW 264.7 whole cell lysate: sc-2211.


RESEARCH USE

For research use only, not for use in diagnostic procedures.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgGκ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

DATA



PBK (11): sc-293028. Western blot analysis of PBK expression in SW480 (A), 3T3-L1 (B), RAW 264.7 (C), F9 (D) and PC-12 (E) whole cell lysates.



PBK (11): sc-293028. Immunoperoxidase staining of formalin fixed, paraffin-embedded human testis tissue showing nuclear staining of cells in seminiferous ducts.

SELECT PRODUCT CITATIONS

- Ohashi, T., et al. 2016. Overexpression of PBK/TOPK contributes to tumor development and poor outcome of esophageal squamous cell carcinoma. *Anticancer Res.* 36: 6457-6466.
- Ohashi, T., et al. 2017. Overexpression of PBK/TOPK relates to tumour malignant potential and poor outcome of gastric carcinoma. *Br. J. Cancer* 116: 218-226.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA