

PAR-4 (5F4): sc-293206

BACKGROUND

Thrombin receptor (also designated protease-activated receptor-1 or PAR-1), PAR-2, PAR-3 and PAR-4 compose a distinct class of G protein-coupled receptors activated by proteolysis. Cleavage of these receptors by proteases occurs within the amino-terminal extracellular domain. Thrombin, a serine protease involved in platelet aggregation and blood coagulation, activates the thrombin receptor, resulting in elevated intracellular calcium levels in platelets. Thrombin also cleaves PAR-3 *in vitro*, suggesting that PAR-3 may be involved in thrombosis or mitogenesis. Thrombin receptor and PAR-4 appear to account for most thrombin signaling in platelets. Activation of PAR-2 *in vitro* is induced by trypsin, suggesting that PAR-2 is not an alternative thrombin receptor. Cytokines including TNF- α and IL-1 β increase PAR-2 expression, indicating PAR-2 involvement in the acute inflammatory response.

REFERENCES

1. Santulli, R.J., et al. 1995. Evidence for the presence of a protease-activated receptor distinct from the thrombin receptor in human keratinocytes. Proc. Natl. Acad. Sci. USA 92: 9151-9155.
2. Lerner, D.J., et al. 1996. Agonist recognition by proteinase-activated receptor 2 and thrombin receptor. Importance of extracellular loop interactions for receptor function. J. Biol. Chem. 271: 13943-13947.
3. Nystedt, S., et al. 1996. The proteinase-activated receptor 2 is induced by inflammatory mediators in human endothelial cells. Comparison with the thrombin receptor. J. Biol. Chem. 271: 14910-14915.
4. Xu, W.F., et al. 1998. Cloning and characterization of human protease-activated receptor 4. Proc. Natl. Acad. Sci. USA 95: 6642-6646.
5. Goldsack, N.R., et al. 1998. Thrombin. Int. J. Biochem. Cell Biol. 30: 641-646.
6. Sullivan, R., et al. 1998. Analysis of a Ca²⁺-activated K⁺ channel that mediates hyperpolarization via the thrombin receptor pathway. Am. J. Physiol. 275: C1342-C1348.
7. Schmidt, V.A., et al. 1998. The human proteinase-activated receptor-3 (PAR-3) gene. Identification within a PAR gene cluster and characterization in vascular endothelial cells and platelets. J. Biol. Chem. 273: 15061-15068.
8. Kahn, M.L., et al. 1999. Protease-activated receptors 1 and 4 mediate activation of human platelets by thrombin. J. Clin. Invest. 103: 879-887.

CHROMOSOMAL LOCATION

Genetic locus: F2RL3 (human) mapping to 19p13.11.

SOURCE

PAR-4 (5F4) is a mouse monoclonal antibody raised against amino acids 18-82 of PAR-4 of human origin.

PRODUCT

Each vial contains 100 μ g IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

PAR-4 (5F4) is recommended for detection of PAR-4 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)] and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


Suitable for use as control antibody for PAR-4 siRNA (h): sc-72068, PAR-4 shRNA Plasmid (h): sc-72068-SH and PAR-4 shRNA (h) Lentiviral Particles: sc-72068-V.

Molecular Weight of PAR-4: 38 kDa.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).

DATA



PAR-4 (5F4): sc-293206. Western blot analysis of human recombinant PAR-4 fusion protein.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.